

Saddle Points in the Auxiliary Field Method

Hiroki Aono and Taro Kashiwa*

Department of Physics, Graduate School of Science and Engineering,
Ehime University, Matsuyama 790-8577, Japan

March 18, 2009

Abstract

Investigations are made on the saddle point calculations (SPC) under the auxiliary field method in path integrations. Two different ways of SPC are considered, Method(I) and Method(II), to be checked in an integral representation of the Gamma function, $\Gamma(N)$, as a bosonic example and in a four-fermi type of Grassmann integral where one "fermion mass" ω_0 differs from the other N -degenerate species. The recipe of Method(I) seems rather complicated than that of (II) superficially, but the case turns out to be opposite in the actual situation. A general formalism allows us to calculate for $\Gamma(N)$ up to $O(1/N^{14})$. It is found that both happen to coincide in the bosonic case but in the fermionic case Method(II) shows a huge deviation in the weak coupling region where $\omega_0 \ll 1$.

*kashiwa@phys.sci.ehime-u.ac.jp

1 Introduction

The auxiliary field method (AFM) is one of the most powerful approximation scheme in path integrations. The recipe for AFM is given as follows: suppose a partition function,

$$Z = \int d^N \boldsymbol{\sigma} d^N \boldsymbol{\sigma}^* \exp \left[-\omega (\boldsymbol{\sigma}^* \cdot \boldsymbol{\sigma}) - \frac{\lambda^2}{2N} (\boldsymbol{\sigma}^* \cdot \boldsymbol{\sigma})^2 \right] ; \quad (\boldsymbol{\sigma}^* \cdot \boldsymbol{\sigma}) \equiv \sum_{a=1}^N \sigma_a^* \sigma_a , \quad (1)$$

where σ_a 's, ($a = 1, \dots, N$), are fermionic or bosonic degrees. ω (called a mass) and λ^2 (a coupling constant) are parameters.

(i) Introduce the auxiliary fields[1, 2], y , (or Hubbard-Stratonovich Field in the solid state physics[3]) by inserting the identity in terms of the Gaussian integral,

$$Z = \int d^N \boldsymbol{\sigma} d^N \boldsymbol{\sigma}^* e^{-\omega (\boldsymbol{\sigma}^* \cdot \boldsymbol{\sigma}) - \frac{\lambda^2}{N} (\boldsymbol{\sigma}^* \cdot \boldsymbol{\sigma})^2} \times \int_{-\infty}^{\infty} \frac{dy}{\sqrt{2\pi}} \exp \left[-\frac{1}{2} \left\{ y + i \frac{\lambda}{\sqrt{N}} (\boldsymbol{\sigma}^* \cdot \boldsymbol{\sigma}) \right\}^2 \right] , \quad (2)$$

in order to remove the four-body interaction, to obtain

$$Z = \int_{-\infty}^{\infty} \frac{dy}{\sqrt{2\pi}} e^{-y^2/2} \int d^N \boldsymbol{\sigma} d^N \boldsymbol{\sigma}^* \exp \left[- \left(\omega + i \frac{\lambda}{\sqrt{N}} y \right) (\boldsymbol{\sigma}^* \cdot \boldsymbol{\sigma}) \right] . \quad (3)$$

(ii) Put $y \mapsto \sqrt{N}y$ and perform the ‘‘Gaussian’’ type integration of σ_a to write

$$E(y) \equiv \int d\sigma_a d\sigma_a^* \exp [- (\omega + i\lambda y) \sigma_a^* \sigma_a] ; \quad \text{for } \forall a . \quad (4)$$

Thus $Z \propto E^N$, yielding

$$Z = \sqrt{\frac{N}{2\pi}} \int_{-\infty}^{\infty} dy \exp \left[-N \left(\frac{y^2}{2} - \ln E(y) \right) \right] . \quad (5)$$

(iii) Write

$$f(y) \equiv \frac{y^2}{2} - \ln E(y) , \quad (6)$$

and assume $N \mapsto \infty$ to perform a saddle point calculation (SPC): find saddle point(s) y_0 , satisfying the stability condition,

$$f'(y) \Big|_{y_0} = 0 , \quad f^{(2)}(y_0) > 0 ; \text{ (stability condition) } , \quad (7)$$

and expand $f(y)$ around y_0 , which gives us a power series of $1/N$, called the loop expansion[4]. This is not a convergent but an asymptotic series, of course.

The prescription is simple and straightforward compared to other nonperturbative methods such as the variational[5] and the optimized perturbation[6]. Moreover, the studies in 1- dimensional(= a quantum mechanical) as well as 0-dimensional(= an integration) bosonic[7] and fermionic[8, 9] models tell us that we can obtain a fairly accurate result, **even when $N = 1$ or small**, from the weak to the strong coupling λ , by taking higher loops into consideration properly.

Now consider a slightly generalized model;

$$Z^{\omega} = \int d^{N+1}\sigma d^{N+1}\sigma^* \exp \left[-(\sigma^* \cdot \omega \cdot \sigma) - \frac{\lambda^2}{2N} (\sigma^* \cdot \sigma)^2 \right] , \quad (8)$$

where $(N+1) \times (N+1)$ matrix ω is given

$$\omega = \begin{pmatrix} \omega_0 & \mathbf{0}^T \\ \mathbf{0} & \omega \mathbf{I}_N \end{pmatrix} , \quad (9)$$

with $\mathbf{I}_N$ and $\mathbf{0}$ being the $N \times N$ unit matrix and the N dimensional zero vector. This corresponds to a field theoretical model of interacting σ_0 and σ_a ($a = 1, 2, \dots, N$) bosons or fermions with the "mass" ω_0 and ω respectively. Introduce an auxiliary field, y , as the above, to find

$$Z^{\omega} = \sqrt{\frac{N}{2\pi}} \int_{-\infty}^{\infty} dy g(y) \exp[-Nf(y)] , \quad (10)$$

where

$$g(y) \equiv \int d\sigma_0 d\sigma_0^* \exp[-(\omega_0 + i\lambda y) \sigma_0^* \sigma_0] , \quad (11)$$

and $f(y)$ has been given by (6), with

$$E(y) \equiv \int d\sigma_a d\sigma_a^* \exp[-(\omega + i\lambda y) \sigma_a^* \sigma_a] ; \quad \text{for } \forall a . \quad (12)$$

In view of (10) an issue comes up: there are two ways of performing SPC.

- Method(I): in¹,

$$I_N \equiv \int dt g(t) e^{-Nf(t)} , \quad (13)$$

find the saddle point t_0 of $f(t)$, $f'(t_0) = 0$, then expand $f(t)$ as well as $g(t)$ around t_0 . Here and hereafter we adopt t instead of y as the integration variable.

- Method(II): rewrite (13) as

$$I_N \equiv \int dt e^{-N\tilde{f}(t)} , \quad \tilde{f}(t) \equiv f(t) - \frac{1}{N} \ln g(t) , \quad (14)$$

then find the saddle t_c of $\tilde{f}(t)$, $\tilde{f}'(t_c) = 0$, and expand $\tilde{f}(t)$ around t_c . Finally put t_c , given in terms of $1/N$ series, into the expression.

If N becomes large both results would match but, as mentioned above, our interest is to study the validity of AFM when N is small. In this paper, therefore, we study the difference between two methods by considering bosonic and fermionic integrations (0-dimensional field theoretical models). In Sec.2, we develop a general formalism of SPC and calculate the asymptotic expansion of the Gamma function as a bosonic model in Sec.3. In Sec.4, we examine a four-fermi type Grassmann integral and the final Sec.5 is devoted to a discussion.

¹The integration range of t , in (13) and (14), need not be specified. Those are given from $-\infty$ to ∞ in the final expression under the saddle point method.(See (15) to (18).)

2 Saddle Points and the Asymptotic Expansion

In this section, we develop a general formalism of SPC.

• Method(I): start with the expression (13) and expand all the integrands around the saddle point, t_0 , to find

$$I_N = e^{-Nf_0} \int dt \left(\sum_{m=0}^{\infty} \frac{g_0^{(m)}}{m!} (t - t_0)^m \right) \exp \left[-N \frac{f_0^{(2)}}{2} (t - t_0)^2 - R_N \right] , \quad (15)$$

where

$$R_N \equiv N \sum_{n=3}^{\infty} \frac{f_0^{(n)}}{n!} (t - t_0)^n , \quad (16)$$

with

$$f_0^{(n)} \equiv f^{(n)}(t_0) , \quad g_0^{(n)} \equiv g^{(n)}(t_0) . \quad (17)$$

After checking the stability condition (7), $f_0^{(2)} > 0$, put $t - t_0 = x / \sqrt{N f_0^{(2)}}$, while assuming N large in (15), to find

$$I_N \cong \frac{e^{-Nf_0}}{\sqrt{N f_0^{(2)}}} \sum_{k,m=0}^{\infty} \frac{(-)^k g_0^{(m)}}{m! k! (N f_0^{(2)})^{m/2}} \int_{-\infty}^{\infty} dx x^m (R_N(x))^k e^{-x^2/2} , \quad (18)$$

where (from (16))

$$R_N(x) \equiv \frac{1}{\sqrt{N}} \sum_{n=0}^{\infty} \frac{F_{n+3}}{(n+3)!} \frac{x^{n+3}}{N^{n/2}} ; \quad F_n \equiv \frac{f_0^{(n)}}{(f_0^{(2)})^{n/2}} . \quad (19)$$

(The integration range of x now stretches from $-\infty$ to ∞ .) Therefore,

$$\begin{aligned} I_N \cong & \frac{e^{-Nf_0}}{\sqrt{N f_0^{(2)}}} \sum_{k,m=0}^{\infty} \frac{(-)^k}{m! k!} \frac{g_0^{(m)}}{(f_0^{(2)})^{m/2}} \sum_{n_1, n_2, \dots, n_k=0}^{\infty} \left(\frac{1}{N} \right)^{(m+k+\sum_{j=1}^k n_j)/2} \\ & \times \frac{F_{n_1+3}}{(n_1+3)!} \frac{F_{n_2+3}}{(n_2+3)!} \cdots \frac{F_{n_k+3}}{(n_k+3)!} \int_{-\infty}^{\infty} dx x^{m+3k+\sum_{j=1}^k n_j} e^{-x^2/2} . \end{aligned} \quad (20)$$

Now put

$$L \equiv \frac{m+k+\sum_{j=1}^k n_j}{2} , \quad (21)$$

which must be integers, $L \in \mathbb{Z}$, not half-integers, $L \in \mathbb{Z} + 1/2$; since the power of x in (20) reads

$$m+3k+\sum_{j=1}^k n_j = 2(L+k) , \quad (22)$$

so that the integral vanishes unless $L + k \in \mathbb{Z}$, leaving us

$$\int_{-\infty}^{\infty} dx x^{2(L+k)} e^{-x^2/2} = \sqrt{2\pi} (2(L+k) - 1)!! . \quad (23)$$

The expression (20) then turns out to be

$$\begin{aligned} I_N \cong & \sqrt{\frac{2\pi}{N f_0^{(2)}}} e^{-N f_0} \sum_{L=0}^{\infty} \frac{1}{N^L} \sum_{k=0}^{2L} \frac{(-)^k}{k!} (2(L+k) - 1)!! \\ & \times \sum_{\substack{\sum_{j=1}^k n_j \leq 2L-k \\ \text{all possible } \{n_j\}}} F(n_1, n_2, \dots, n_k) , \end{aligned} \quad (24)$$

where

$$F(n_1, n_2, \dots, n_k) \equiv \frac{1}{(2L - k - \sum_{j=1}^k n_j)!} \frac{g_0^{(2L-k-\sum_{j=1}^k n_j)}}{(f_0^{(2)})^{(2L-k-\sum_{j=1}^k n_j)/2}} \prod_{j=1}^k \frac{F_{n_j+3}}{(n_j+3)!} , \quad (25)$$

is a symmetric function of n_j 's. In (24) the sum should be taken under the condition,

$$\sum_{j=1}^k n_j \leq 2L - k ; \quad (26)$$

since $m \geq 0$ in (22). Note if $k = 0$, then $\sum_{j=1}^0 n_j \equiv 0$ and $\prod_{j=1}^0 G(n_j) \equiv 1$, so that

$$F(n_1, \dots, n_k) \stackrel{k=0}{=} \frac{1}{(2L)!} \frac{g_0^{(2L)}}{(f_0^{(2)})^L} . \quad (27)$$

The conditional sum of n_j 's, (26), can be expressed as an alternative form: suppose Q_α of n_j 's are alike of A_α , $n_{j_1} = n_{j_2} = \dots = n_{j_{Q_\alpha}} = A_\alpha$, to write

$$F \left(\overbrace{A_1, \dots, A_1}^{Q_1}, \overbrace{A_2, \dots, A_2}^{Q_2}, \dots, \overbrace{A_P, \dots, A_P}^{Q_P} \right) \equiv F \left(\{A_1^{Q_1}\}, \{A_2^{Q_2}\}, \dots, \{A_P^{Q_P}\} \right) . \quad (28)$$

It is clear that the condition (26) reads as

$$P \leq k ; \quad \sum_{\alpha=1}^P Q_\alpha = k , \quad \sum_{\alpha=1}^P Q_\alpha A_\alpha \left(= \sum_{j=1}^k n_j \right) \leq 2L - k . \quad (29)$$

The multiplicity reads

$$\binom{k}{Q_1} \binom{k-Q_1}{Q_2} \dots \binom{k-\sum_{\alpha=1}^{P-1} Q_\alpha}{Q_P} = \frac{k!}{Q_1! \dots Q_P!} , \quad (30)$$

so that (24) becomes

$$I_N \cong \sqrt{\frac{2\pi}{Nf_0^{(2)}}} e^{-Nf_0} \sum_{L=0}^{\infty} \frac{1}{N^L} \sum_{k=0}^{2L} (-)^k (2(L+k)-1)!! \\ \times \sum_{\substack{\sum_{\alpha=1}^P Q_{\alpha} A_{\alpha} \leq 2L-k \\ \text{all possible } \{A_{\alpha}\} \\ \sum_{\alpha=1}^P Q_{\alpha} = k ; (P \leq k)}} \frac{1}{Q_1! \cdots Q_P!} F\left(\{A_1^{Q_1}\}, \{A_2^{Q_2}\}, \dots, \{A_P^{Q_P}\}\right) , \quad (31)$$

where we have assumed that $A_1 < A_2 < \cdots < A_P$. This is the main formula of the Method(I). Let us classify I_N , according to the WKB-approximation [4] under which $1/\hbar$ appears instead of N ;

1. Tree:

$$(I_N)_{\text{tree}} \equiv e^{-Nf_0} g_0 . \quad (32)$$

2. l -loop: terms up to $l (\geq 1)$ in (24):

$$(I_N)_{l\text{-loop}} \equiv e^{-Nf_0} \sqrt{\frac{2\pi}{Nf_0^{(2)}}} g_0 \left(1 + \cdots + O\left(\frac{1}{N^{l-1}}\right) \right) . \quad (33)$$

• Method(II): by putting $f(t) \mapsto \tilde{f}(t)$; $g(t) \mapsto 1$ and $t_0 \mapsto t_c$, all expressions in Method(I) can be read as those of Method(II). Write

$$\tilde{f}_c^{(n)} \equiv \tilde{f}^{(n)}(t_c) , \quad (34)$$

then (24) is changed to

$$I_N \cong \sqrt{\frac{2\pi}{N\tilde{f}_c^{(2)}}} e^{-N\tilde{f}_c} \sum_{L=0}^{\infty} \frac{1}{N^L} \\ \times \sum_{k=0}^{2L} \frac{(-)^k}{k!} (2(L+k)-1)!! \sum_{\substack{\sum_{j=1}^k n_j = 2L-k \\ \text{all possible } \{n_j\}}} \tilde{F}(n_1, n_2, \dots, n_k) , \quad (35)$$

$$\tilde{F}(n_1, n_2, \dots, n_k) \equiv \prod_{j=1}^k \frac{\tilde{F}_{n_j+3}}{(n_j+3)!} ; \quad \tilde{F}_n \equiv \frac{\tilde{f}_c^{(n)}}{(\tilde{f}_c^{(2)})^{n/2}} . \quad (36)$$

It should be noted that without $g(t)$ the conditional sum is given **by the equality**, $\sum_{j=1}^k n_j = 2L - k$, **not by the inequality** (26). Also (31) is changed to

$$I_N \cong \sqrt{\frac{2\pi}{N\tilde{f}_c^{(2)}}} e^{-N\tilde{f}_c} \sum_{L=0}^{\infty} \frac{1}{N^L} \sum_{k=0}^{2L} (-)^k (2(L+k)-1)!! \\ \times \sum_{\substack{\sum_{\alpha=1}^P Q_{\alpha} A_{\alpha} = 2L-k \\ \text{all possible } \{A_{\alpha}\} \\ \sum_{\alpha=1}^P Q_{\alpha} = k ; (P \leq k)}} \frac{1}{Q_1! \cdots Q_P!} \tilde{F}\left(\{A_1^{Q_1}\}, \{A_2^{Q_2}\}, \dots, \{A_P^{Q_P}\}\right) , \quad (37)$$

where $\tilde{F}\left(\{A_1^{Q_1}\}, \{A_2^{Q_2}\}, \dots, \{A_P^{Q_P}\}\right)$ is defined by the expression, (28), with the tildes. Here again the conditional sum (29) becomes simpler, $\sum_{\alpha=1}^P Q_{\alpha} A_{\alpha} = 2L - k$. Classification in this case is again

1. Tree:

$$(I_N)_{\text{tree}} \equiv e^{-N\tilde{f}_c^{(0)}} , \quad (38)$$

2. l -loop: terms up to $l (\geq 1)$ in (20) or (24):

$$(I_N)_{l\text{-loop}} \equiv e^{-N\tilde{f}_c^{(0)}} \sqrt{\frac{2\pi}{N\tilde{f}_c^{(2)}}} \left(1 + \dots + O\left(\frac{1}{N^{l-1}}\right) \right) . \quad (39)$$

Note that there are additional powers of $1/N$ hidden in t_c . The final task is then **to expand all functions of t_c up to $(1/N)^{l-1}$** , which however depends on individual models so is relegated to the following sections.

3 A Bosonic Case: the Gamma Function

As a simple bosonic example, in this section we consider the Gamma function,

$$\Gamma(N) = \int_0^\infty dt t^{N-1} e^{-t} = N^N I_N ; \quad I_N \equiv \int_0^\infty dt \left(\frac{1}{t} \right) e^{-N(t - \ln t)} , \quad (40)$$

where we have put $t \mapsto Nt$ in the final expression. Then

$$\text{Method(I)} : \quad f(t) \equiv t - \ln t ; \quad g(t) \equiv \frac{1}{t} . \quad (41)$$

$$\text{Method(II)} : \quad \tilde{f}(t) \equiv t - \left(1 - \frac{1}{N} \right) \ln t . \quad (42)$$

•Method(I): the saddle point, t_0 is given by

$$f'(t_0) = 1 - \frac{1}{t_0} = 0 \implies t_0 = 1 , \quad (43)$$

so that

$$f_0 = 1 = g_0 , \quad f_0^{(n+1)} = -g_0^{(n)} = -(-)^n n! ; \quad (n \geq 1) , \quad (44)$$

which ensures the stability condition, $f_0^{(2)} = 1 > 0$, and

$$F(n_1, n_2, \dots, n_k) = \prod_{j=1}^k \frac{1}{(n_j + 3)} , \quad (45)$$

from (25). Therefore from (24) and (31),

$$I_N \cong \sqrt{\frac{2\pi}{N}} e^{-N} \sum_{L=0}^{\infty} \frac{1}{N^L} \sum_{k=0}^{2L} T(L, k | \leq 2L - k) , \quad (46)$$

where

$$\begin{aligned}
T(L, k | \leq 2L - k) &\equiv \frac{(-)^k}{k!} (2(L + k) - 1)!! \sum_{\substack{\sum_{j=1}^k n_j \leq 2L - k \\ \text{all possible } \{n_j\}}} \prod_{j=1}^k \frac{1}{(n_j + 3)} \quad (47) \\
&= (-)^k (2(L + k) - 1)!! \sum_{\substack{\sum_{\alpha=1}^P Q_\alpha A_\alpha \leq 2L - k \\ \text{all possible } \{A_\alpha\} \\ \sum_{\alpha=1}^P Q_\alpha = k ; (P \leq k)}} \frac{1}{Q_1! \cdots Q_P!} \\
&\quad \times \frac{1}{(A_1 + 3)^{Q_1} (A_2 + 3)^{Q_2} \cdots (A_P + 3)^{Q_P}} . \quad (48)
\end{aligned}$$

Here we concentrate on the latter expression (48) and realize that most of the terms in the conditional sum, $\sum_{\alpha=1}^P Q_\alpha A_\alpha \leq 2L - k$, are canceled, leaving only the terms satisfying the equality $\sum_{\alpha=1}^P Q_\alpha A_\alpha = 2L - k$, that is

$$\sum_{k=0}^{2L} T(L, k | \leq 2L - k) = \sum_{k=0}^{2L} T(L, k | 2L - k) , \quad (49)$$

to give

$$I_N \cong \sqrt{\frac{2\pi}{N}} e^{-N} \sum_{L=0}^{\infty} \frac{1}{N^L} \sum_{k=0}^{2L} T(L, k | 2L - k) , \quad (50)$$

where

$$\begin{aligned}
T(L, k | 2L - k) &\equiv \frac{(-)^k}{k!} (2(L + k) - 1)!! \sum_{\substack{\sum_{j=1}^k n_j = 2L - k \\ \text{all possible } \{n_j\}}} \prod_{j=1}^k \frac{1}{(n_j + 3)} \quad (51) \\
&= (-)^k (2(L + k) - 1)!! \sum_{\substack{\sum_{\alpha=1}^P Q_\alpha A_\alpha = 2L - k \\ \text{all possible } \{A_\alpha\} \\ \sum_{\alpha=1}^P Q_\alpha = k ; (P \leq k)}} \frac{1}{Q_1! \cdots Q_P!} \\
&\quad \times \frac{1}{(A_1 + 3)^{Q_1} (A_2 + 3)^{Q_2} \cdots (A_P + 3)^{Q_P}} , \quad (52)
\end{aligned}$$

with the definition when $k = 0$, (implying $P = 0$), and $L \neq 0$

$$T(L, 0 | 2L) = 0 , \quad (53)$$

since otherwise there is inconsistency in the conditional sum $\sum_{\alpha=1}^{P=0} Q_\alpha A_\alpha (= 0)$ and $2L (\neq 0)$. The numerical values of $T(L, k | 2L - k)$ (apart from the factor $(2(L + k) - 1)!!$) are listed in the appendix A. The proof of (49) is rather lengthy then relegated to the appendix B.

In view of (24) and (25), the above fact implies that there is no contribution from $g_0^{(m)} (m \geq 1)$, that is, the equality holds

$$\int_0^\infty dt \left(\frac{1}{t} \right) e^{-N(t - \ln t)} \stackrel{1/N}{=} \int_0^\infty dt e^{-N(t - \ln t)} , \quad (54)$$

under the $1/N$ expansion (Actually it does hold without $1/N$ in this case: see the expression (167). A detailed discussion is relegated to the appendix B.)

In view of (32) as well as (40),

$$(\Gamma(N))_{\text{tree}} = N^N e^{-N} . \quad (55)$$

And the result of 15-loop is from (33),

$$\begin{aligned} (\Gamma(N))_{15\text{-loop}} = & \left(\sqrt{\frac{2\pi}{N}} N^N e^{-N} \right) \left\{ 1 + \frac{1}{12} \frac{1}{N} + \frac{1}{288} \frac{1}{N^2} - \frac{139}{51840} \frac{1}{N^3} - \frac{571}{2488320} \frac{1}{N^4} \right. \\ & + \frac{163879}{209018880} \frac{1}{N^5} + \frac{5246819}{75246796800} \frac{1}{N^6} - \frac{534703531}{902961561600} \frac{1}{N^7} \\ & - \frac{4483131259}{86684309913600} \frac{1}{N^8} + \frac{432261921612371}{514904800886784000} \frac{1}{N^9} \\ & + \frac{6232523202521089}{86504006548979712000} \frac{1}{N^{10}} - \frac{13494625021640835072000}{1579029138854919086429} \frac{1}{N^{11}} \\ & - \frac{1579029138854919086429}{9716130015581401251840000} \frac{1}{N^{12}} + \frac{746590869962651602203151}{116593560186976815022080000} \frac{1}{N^{13}} \\ & \left. + \frac{1511513601028097903631961}{2798245444487443560529920000} \frac{1}{N^{14}} + O(N^{-15}) \right\} . \quad (56) \end{aligned}$$

This coincides the results in references²; up to $1/N^4$ [10], $1/N^7$ [11] and $1/N^9$ [12].




Figure 1: The ratio of Approximation to Exact in Method(I): the horizontal axis designates L , $0 \leq L \leq 14$, omitting the tree part. The solid line with the circle, the dotted with the triangle, the dashed with the cross, and the dash-dotted with the square designate $N = 1, 2, 5$ and 10 respectively. The tail end of the $N = 1$ line deviates from the unity, implying the asymptotic character of the $1/N$ expansion.

In the table 1, we list $N = 1, 2, 5$, and 10 results up to 15-loop including the tree ones, where the optimized values are shaded. Also we plot the ratio of approximate to exact values for $0 \leq L \leq 14$ (omitting the tree part) in the figure 1. From these we convince the validity of the loop expansion in this case; since even in the smallest $N = 1$ case, the 2- or 3-loop approximation gives ~ 0.1 or $\sim 0.2\%$ error. It also should be noted that the characteristic feature of the asymptotic expansion can be read from deviation after passing through the optimized values, which is most clearly seen in the figure 1 at the tail end of the $N = 1$ line.

²We have no information how those results are obtained, but the derivations would be from another prescription such as the Watson's Lemma[13]

	$N = 1$	$N = 2$	$N = 5$	$N = 10$
Exact	1	1	24	362880
tree	0.36788	0.54134	2.10561×10^1	4.53999×10^5
(tree)/Exact	0.36788	0.54134	0.87734	1.25110
1-loop	0.92214	0.95950	2.36038×10^1	3.59870×10^5
(1-loop)/Exact	0.92214	0.95950	0.98349	0.99170
2-loop	0.99898	0.99948	2.39972×10^1	3.62868×10^5
(2-loop)/Exact	0.99898	0.99948	0.99988	0.99997
3-loop	1.00218	1.00031	2.40005×10^1	3.62881×10^5
(3-loop)/Exact	1.00218	1.00031	1.00002	1.00000
4-loop	0.99971	0.99999	2.40000×10^1	3.62880×10^5
(4-loop)/Exact	0.99971	0.99999	1.00000	1.00000
5-loop	0.99950	0.99998	2.40000×10^1	3.62880×10^5
(5-loop)/Exact	0.99950	0.99998	1.00000	1.00000
6-loop	1.00022	1.00000	2.40000×10^1	3.62880×10^5
(6-loop)/Exact	1.00022	1.00000	1.00000	1.00000
7-loop	1.00029	1.00000	2.40000×10^1	3.62880×10^5
(7-loop)/Exact	1.00029	1.00000	1.00000	1.00000
8-loop	0.99974	1.00000	2.40000×10^1	3.62880×10^5
(8-loop)/Exact	0.99974	1.00000	1.00000	1.00000
9-loop	0.99969	1.00000	2.40000×10^1	3.62880×10^5
(9-loop)/Exact	0.99969	1.00000	1.00000	1.00000
10-loop	1.00047	1.00000	2.40000×10^1	3.62880×10^5
(10-loop)/Exact	1.00047	1.00000	1.00000	1.00000
11-loop	1.00053	1.00000	2.40000×10^1	3.62880×10^5
(11-loop)/Exact	1.00053	1.00000	1.00000	1.00000
12-loop	0.99877	1.00000	2.40000×10^1	3.62880×10^5
(12-loop)/Exact	0.99877	1.00000	1.00000	1.00000
13-loop	0.99862	1.00000	2.40000×10^1	3.62880×10^5
(13-loop)/Exact	0.99862	1.00000	1.00000	1.00000
14-loop	1.00452	1.00000	2.40000×10^1	3.62880×10^5
(14-loop)/Exact	1.00452	1.00000	1.00000	1.00000
15-loop	1.00502	1.00000	2.40000×10^1	3.62880×10^5
(15-loop)/Exact	1.00502	1.00000	1.00000	1.00000

Table 1: Results of Method(I) in $N = 1, 2, 5$ and 10 up to $L = 14$. Even in $N = 1$, 2-loop($L = 1$) approximation is sufficiently close to the exact value, however, deviation becomes gradually eminent in the loops larger than 7. The optimized values are shaded in each N , whose position shows that the $1/N$ expansion is indeed an asymptotic one.

•Method(II): from (42), the saddle point is

$$\tilde{f}'(t) = 1 - \left(1 - \frac{1}{N}\right) \frac{1}{t} = 0 \implies t_c = t_0 + \frac{t_1}{N} ; \quad t_0 \equiv 1 , \quad t_1 \equiv -1 , \quad (57)$$

and satisfies the stability condition,

$$\tilde{f}_c^{(2)} = \frac{1}{t_c} = \frac{N}{N-1} > 0 . \quad (58)$$

Note that

$$\tilde{f}_c = t_c (1 - \ln t_c) , \quad \tilde{f}_c^{(n)} \equiv \tilde{f}^{(n)}(t_c) = (-)^n (n-1)! \frac{1}{(t_c)^{n-1}} ; \quad (n \geq 2) , \quad (59)$$

to give $\tilde{F}_n = \frac{(-)^n (n-1)!}{t_c^{n/2-1}}$ so that

$$\tilde{F}(n_1, n_2, \dots, n_k) = (-)^{3k + \sum_j n_j} \frac{1}{t_c^{(\sum_j n_j + k)/2}} \prod_{j=1}^k \frac{1}{(n_j + 3)} = \frac{1}{(t_c)^L} \prod_{j=1}^k \frac{1}{(n_j + 3)} , \quad (60)$$

with $\sum_{j=1}^k n_j + k = 2L$, from (36). Therefore with the aid of (35)

$$I_N \cong e^{-N t_c (1 - \ln t_c)} \sqrt{\frac{2\pi t_c}{N}} \sum_{L=0}^{\infty} \frac{1}{N^L} (t_c)^{-L} \sum_{k=0}^{2L} T(L, k | 2L - k) . \quad (61)$$

The tree and the 1-loop part are given

$$(I_N)_{\text{tree}} = e^{-Nt_c(1-\ln t_c)} \Big|_{t_1=0} = e^{-N} , \quad (62)$$

$$(I_N)_{1\text{-loop}} = e^{-Nt_c(1-\ln t_c)} \sqrt{\frac{2\pi t_c}{N}} \Big|_{t_1=0} = e^{-N} \sqrt{\frac{2\pi}{N}} . \quad (63)$$

From the 2-loop approximation, $t_1(57)$ should be included to t_c , and all function of t_c must be expanded up to $O(1/N^L) : L = 1, 2, \dots$ in (61), introduce the prefactor $\mathcal{P}$,

$$\mathcal{P} \equiv e^{-Nt_c(1-\ln t_c)} \sqrt{\frac{2\pi t_c}{N}} = e^{-N} \sqrt{\frac{2\pi}{N}} \left[1 - \frac{1}{12} \frac{1}{N^2} - \frac{1}{12} \frac{1}{N^3} - \frac{103}{1440} \frac{1}{N^4} + O\left(\frac{1}{N^5}\right) \right] , \quad (64)$$

and the loop factor $\mathcal{L}$

$$\begin{aligned} \mathcal{L} \equiv \sum_{L=0}^{\infty} \frac{1}{N^L} (t_c)^{-L} \sum_{k=0}^{2L} T(L, k|2L-k) &= 1 + \frac{n(1)}{N} + \frac{n(1)+n(2)}{N^2} \\ &+ \frac{n(1)+2n(2)+n(3)}{N^3} + \frac{n(1)+3n(2)+3n(3)+n(4)}{N^4} + O\left(\frac{1}{N^5}\right) , \end{aligned} \quad (65)$$

where

$$n(L) \equiv \sum_{k=0}^{2L} T(L, k|2L-k) ; \quad (66)$$

$$n(1) = \frac{1}{12} , \quad n(2) = \frac{1}{288} , \quad n(3) = -\frac{139}{51840} , \quad n(4) = -\frac{571}{2488320} , \quad (67)$$

from the table 5 in the appendix A. $\mathcal{P} \times \mathcal{L}$ gives the 5-loop approximation of I_N

$$(I_N)_{5\text{-loop}} = e^{-N} \sqrt{\frac{2\pi}{N}} \left[1 + \sum_{L=1}^4 \frac{n(L)}{N^L} \right] , \quad (68)$$

which is exactly the same to the one ($L \mapsto 4$ in (50)) in Method(I). Many terms in the numerator, $n(1), \dots, n(L-1)$ are canceled, leaving us only $n(L)$. These miracle cancellations occur for all orders of $1/N$, yielding the result that **there is no difference between Method(I) and (II)** in this case.

The reason is rather easily figured out: because of the formula (167) in the appendix B, we can put $g(t) \equiv 1/t$ in the integral I_N (40) to the unity, $g(t) \mapsto 1$. In other words, **Method(I) is equivalent to Method(II)** in the Gamma function case.

4 A Fermionic Case

Our target is a fermionic version of (8);

$$Z \equiv \int d^{N+1} \hat{\xi} d^{N+1} \hat{\xi}^* \exp \left[-\hat{\xi}^* \cdot \omega \cdot \hat{\xi} + \frac{\lambda^2}{2N} (\hat{\xi}^* \cdot \hat{\xi})^2 \right] , \quad (69)$$

where $\hat{\xi}, \hat{\xi}^*(\xi, \xi^*)$ are $N+1(N)$ -dimensional Grassmann variables,

$$\hat{\xi} \equiv (\xi_0, \xi) , \quad \hat{\xi}^* \equiv (\xi_0^*, \xi^*) ; \quad \xi \equiv (\xi_1, \dots, \xi_N) , \quad \xi^* \equiv (\xi_1^*, \dots, \xi_N^*) , \quad (70)$$

with

$$d^{N+1}\hat{\xi} \equiv d^N \xi d\xi_0 \equiv d\xi_N d\xi_{N-1} \cdots d\xi_0 , \quad d^{N+1}\hat{\xi}^* \equiv d\xi_0^* d^N \xi^* \equiv d\xi_0^* d\xi_1^* \cdots d\xi_N^* , \quad (71)$$

and $(N+1) \times (N+1)$ matrix, ω , has been given in (9). Z is calculable by means of a standard Grassmann integration;

$$\int d^n \xi d^n \xi^* (\xi^* \cdot \xi)^m = (-)^n n! \delta_{mn} , \quad (72)$$

to obtain

$$Z = \sum_{r=0}^{\lfloor \frac{N}{2} \rfloor} \frac{N!}{r!(N-2r)!} \left(\omega_0 \omega + \frac{\lambda^2}{N} (N-2r) \right) (\omega)^{N-2r-1} \left(\frac{\lambda^2}{2N} \right)^r . \quad (73)$$

In this analysis, we assume that all parameters in this model are real and positive³, $\omega > 0, \omega_0 > 0, \lambda > 0$, and take $N = 2$ with

$$0 \leq \lambda \leq 10 ; \quad \omega_0 = 10^2 \omega , \quad \omega , \quad 10^{-2} \omega . \quad (74)$$

(The case, $\omega_0 = 10^2 \omega$ is a toy model of u-, d-, and s-quarks.)

Introducing an auxiliary field, in terms of

$$1 = \int_{-\infty}^{\infty} \frac{dy}{\sqrt{2\pi}} \exp \left[-\frac{1}{2} \left(y + \frac{\lambda}{\sqrt{N}} (\hat{\xi}^* \cdot \hat{\xi}) \right)^2 \right] , \quad (75)$$

into the target (69), we obtain

$$Z = \sqrt{\frac{N}{2\pi}} \int_{-\infty}^{\infty} dy (\omega_0 + \lambda y) \exp \left[-N \left(\frac{y^2}{2} - \ln(\omega + \lambda y) \right) \right] , \quad (76)$$

where we have performed the Grassmann Gaussian integration,

$$\int d\xi d\xi^* e^{-\omega \xi^* \xi} = \omega , \quad (77)$$

and y has been scaled, $y \mapsto \sqrt{N}y$, as before. Now write

$$Z = \sqrt{\frac{N}{2\pi}} I_N ; \quad I_N \equiv \int dt g(t) e^{-Nf(t)} , \quad (78)$$

$$g(t) \equiv \omega_0 + \lambda t ; \quad f(t) \equiv \frac{t^2}{2} - \ln(\omega + \lambda t) , \quad (79)$$

for Method(I) and

$$I_N = \int dt e^{-N\tilde{f}(t)} ,$$

$$\tilde{f}(t) \equiv f(t) - \frac{1}{N} \ln g(t) = \frac{t^2}{2} - \ln(\omega + \lambda t) - \frac{1}{N} \ln(\omega_0 + \lambda t) , \quad (80)$$

for Method(II). (Here y has been switched to t .)

³Although we can see an interesting phenomenon when $\lambda^2 < 0$:the caustics emerge[9].

•Method(I): the saddle points are determined by

$$f^{(1)}(t)\Big|_{t=t_0} = t_0 - \frac{\lambda}{\omega + \lambda t_0} = 0 . \quad (81)$$

Here and hereafter the equation is called as the gap equation[14]. If we introduce

$$\Omega_0 \equiv \omega + \lambda t_0 , \quad (82)$$

(81) becomes

$$(\Omega_0)^2 - \omega \Omega_0 - \lambda^2 = 0 , \quad (83)$$

yielding to two saddle points

$$\Omega_0^{(\pm)} \equiv \frac{\omega \pm \sqrt{\omega^2 + 4\lambda^2}}{2} . \quad (84)$$

The stability condition (7) in this case reads

$$f_0^{(2)} = 1 + \left(\frac{\lambda}{\Omega_0} \right)^2 = \frac{(\Omega_0)^2 + \lambda^2}{(\Omega_0)^2} > 0 ; \quad (85)$$

which is positive for both $\Omega_0^{(\pm)}$. The value of $f(t)$, $g(t)$ and derivatives at the saddle points are given by

$$f_0 = \frac{(\Omega_0 - \omega)^2}{2\lambda^2} - \ln \Omega_0 \quad ; \quad f_0^{(m)} = (m-1)! \left(-\frac{\lambda}{\Omega_0} \right)^m \quad ; \quad (m \geq 3) , \quad (86)$$

$$g_0 = \Omega_0 + \delta\omega \quad ; \quad g_0^{(1)} = \lambda \quad ; \quad g_0^{(m)} = 0 \quad ; \quad (m \geq 2) , \quad (87)$$

where

$$\delta\omega \equiv \omega_0 - \omega . \quad (88)$$

Then from (19)

$$\frac{F_{n_j+3}}{(n_j+3)!} = \frac{1}{(n_j+3)!} \left(f_0^{(2)} \right)^{-(n_j+3)/2} f_0^{(n_j+3)} = \frac{1}{n_j+3} \left(\frac{-\epsilon(\Omega_0) \lambda}{\sqrt{(\Omega_0)^2 + \lambda^2}} \right)^{n_j+3} , \quad (89)$$

where

$$\epsilon(\Omega_0) = \begin{cases} 1 & : \quad \Omega_0 > 0 \\ -1 & : \quad \Omega_0 < 0 \end{cases} , \quad (90)$$

is the sign function. Accordingly, in view of (24) with (25), we find

$$\begin{aligned} & \frac{(-)^k}{k!} (2(L+k)-1)!! \sum_{\substack{\sum_{j=1}^k n_j \leq 2L-k \\ \text{all possible } \{n_j\}}} F(n_1, n_2, \dots, n_k) \\ &= \left(\frac{\lambda}{\sqrt{(\Omega_0)^2 + \lambda^2}} \right)^{2(L+k)} [(\Omega_0 + \delta\omega) T(L, k|2L-k) - \Omega_0 T(L, k|2L-k-1)] , \quad (91) \end{aligned}$$

where use has been made of the notation (51): $T(L, k|2L - k - 1)$ is defined by replacing the sum $\sum_{j=1}^k n_j = 2L - k$ to $\sum_{j=1}^k n_j = 2L - k - 1$ in (51). Therefore from (25)

$$I_N \cong \sqrt{\frac{2\pi}{N}} \epsilon(\Omega_0) \frac{\Omega_0^{N+1}}{\sqrt{(\Omega_0)^2 + \lambda^2}} \exp \left[-N \frac{(\Omega_0 - \omega)^2}{2\lambda^2} \right] \sum_{L=0}^{\infty} \frac{1}{N^L} \sum_{k=0}^{2L} \left(\frac{\lambda}{\sqrt{(\Omega_0)^2 + \lambda^2}} \right)^{2(L+k)} \times [(\Omega_0 + \delta\omega) T(L, k|2L - k) - \Omega_0 T(L, k|2L - k - 1)] , \quad (92)$$

so that

$$Z \cong \epsilon(\Omega_0) \frac{\Omega_0^{N+1}}{\sqrt{(\Omega_0)^2 + \lambda^2}} \exp \left[-N \frac{(\Omega_0 - \omega)^2}{2\lambda^2} \right] \sum_{L=0}^{\infty} \frac{1}{N^L} \sum_{k=0}^{2L} \left(\frac{\lambda}{\sqrt{(\Omega_0)^2 + \lambda^2}} \right)^{2(L+k)} \times [(\Omega_0 + \delta\omega) T(L, k|2L - k) - \Omega_0 T(L, k|2L - k - 1)] . \quad (93)$$

According to classification in sec.2, (32) and (33), the tree and the l -loop approximation read

$$Z_{\text{tree}} \equiv e^{-N f_0} g_0 = \exp \left[-N \frac{(\Omega_0 - \omega)^2}{2\lambda^2} \right] \Omega_0^N (\Omega_0 + \delta\omega) , \quad (94)$$

$$Z_{l\text{-loop}} \equiv \epsilon(\Omega_0) \frac{\Omega_0^{N+1}}{\sqrt{(\Omega_0)^2 + \lambda^2}} \exp \left[-N \frac{(\Omega_0 - \omega)^2}{2\lambda^2} \right] \sum_{L=0}^{l-1} \frac{1}{N^L} \sum_{k=0}^{2L} \left(\frac{\lambda}{\sqrt{(\Omega_0)^2 + \lambda^2}} \right)^{2(L+k)} \times [(\Omega_0 + \delta\omega) T(L, k|2L - k) - \Omega_0 T(L, k|2L - k - 1)] . \quad (95)$$

Up to 3-loop ($L \leq 2$), by noting the table 5 in the appendix A and $T(1, 1|0) = -1, T(2, 1|2) = -3, T(2, 2|1) = 35/4, T(2, 3|1) = -35/6$ for $T(L, k|2L - k - 1)$, we find

$$Z_{1\text{-loop}} = \epsilon(\Omega_0) \exp \left[-N \frac{(\Omega_0 - \omega)^2}{2\lambda^2} \right] \frac{(\Omega_0)^{N+1} (\Omega_0 + \delta\omega)}{\sqrt{(\Omega_0)^2 + \lambda^2}} , \quad (96)$$

$$Z_{2\text{-loop}} = Z_{1\text{-loop}} + \epsilon(\Omega_0) \exp \left[-N \frac{(\Omega_0 - \omega)^2}{2\lambda^2} \right] \frac{(\Omega_0)^{N+1}}{\sqrt{(\Omega_0)^2 + \lambda^2}} \times \frac{1}{N} \left[\frac{\lambda^4 (\Omega_0 - 3\delta\omega)}{4 [(\Omega_0)^2 + \lambda^2]^2} + \frac{5\lambda^6 (\Omega_0 + \delta\omega)}{6 [(\Omega_0)^2 + \lambda^2]^3} \right] , \quad (97)$$

and

$$Z_{3\text{-loop}} = Z_{2\text{-loop}} + \epsilon(\Omega_0) \exp \left[-N \frac{(\Omega_0 - \omega)^2}{2\lambda^2} \right] \frac{(\Omega_0)^{N+1}}{\sqrt{(\Omega_0)^2 + \lambda^2}} \frac{1}{N^2} \left[\frac{\lambda^6 (\Omega_0 - 5\delta\omega)}{2 [(\Omega_0)^2 + \lambda^2]^3} + \frac{7\lambda^8 (7\Omega_0 + 47\delta\omega)}{32 [(\Omega_0)^2 + \lambda^2]^4} - \frac{35\lambda^{10} (5\Omega_0 + 9\delta\omega)}{24 [(\Omega_0)^2 + \lambda^2]^5} + \frac{385\lambda^{12} (\Omega_0 + \delta\omega)}{72 [(\Omega_0)^2 + \lambda^2]^6} \right] . \quad (98)$$

Since there are two saddle points $\Omega_0^\pm$ (84) the total Z is given by

$$Z_{l\text{-loop}}^{\text{Total}} = Z_{l\text{-loop}}^{(+)} + Z_{l\text{-loop}}^{(-)} , \quad (99)$$

where $Z_{l\text{-loop}}^{(\pm)}$ has been obtained by putting $\Omega_0 \mapsto \Omega_0^{(\pm)}$ in (95).

In the table 2, we list the result of $\omega_0 = 10^2\omega, \omega, 10^{-2}\omega$ for $10^{-3} \leq \lambda \leq 10$ in $N = 2$. We put $\omega \mapsto 1$ and write the data of the ratio of Z_{tree} and $Z_{l\text{-loop}}(l = 1, 2, 3)$ to the exact value. From this, in the weak coupling region, $\lambda < 1$, the 1-loop approximation almost yields the exact value; even in the worst case, $\omega_0 = 10^{-2}$, only 0.3% error crops up. For a whole coupling region including $\lambda \geq 1$, the error is within 1.1% under 2-loop and becomes venial, $< 0.3\%$, under the 3-loop approximation.

$\omega_0 = 10^2$	λ	Exact	tree (tree)/Exact	1-loop (1-loop)/Exact	2-loop (2-loop)/Exact	3-loop (3-loop)/Exact
	10^{-3}	1.0000×10^2	1.0000×10^2 1.0000	1.0000×10^2 1.0000	1.0000×10^2 1.0000	1.0000×10^2 1.0000
	10^{-2}	1.0001×10^2	1.0001×10^2 1.0000	1.0001×10^2 1.0000	1.0001×10^2 1.0000	1.0001×10^2 1.0000
	10^{-1}	1.0051×10^2	1.0101×10^2 1.0049	1.0051×10^2 1.0000	1.0051×10^2 1.0000	1.0051×10^2 1.0000
	1	1.5100×10^2	1.8253×10^2 1.2088	1.5438×10^2 1.0224	1.5138×10^2 1.0025	1.5096×10^2 0.9998
	10	5.2000×10^3	7.5782×10^3 1.4574	5.3961×10^3 1.0377	5.1861×10^3 0.9973	5.1906×10^3 0.9982
$\omega_0 = \omega$	λ	Exact	tree (tree)/Exact	1-loop (1-loop)/Exact	2-loop (2-loop)/Exact	3-loop (3-loop)/Exact
	10^{-3}	1.0000	1.0000 1.0000	1.0000 1.0000	1.0000 1.0000	1.0000 1.0000
	10^{-2}	1.0002	1.0002 1.0000	1.0001 1.0000	1.0001 1.0000	1.0002 1.0000
	10^{-1}	1.0150	1.0200 1.0049	1.0150 1.0000	1.0150 1.0000	1.0150 1.0000
	1	2.5000	2.8740 1.1496	2.4503 0.9801	2.4934 0.9974	2.4998 0.9999
	10	1.5100×10^2	1.8498×10^2 1.2250	1.4410×10^2 0.9543	1.5030×10^2 0.9954	1.5124×10^2 1.0016
$\omega_0 = 10^{-2}$	λ	Exact	tree (tree)/Exact	1-loop (1-loop)/Exact	2-loop (2-loop)/Exact	3-loop (3-loop)/Exact
	10^{-3}	1.0001×10^{-2}	1.0001×10^{-2} 1.0000	1.0001×10^{-2} 1.0000	1.0001×10^{-2} 1.0000	1.0001×10^{-2} 1.0000
	10^{-2}	1.0101×10^{-2}	1.0101×10^{-2} 1.0000	1.0100×10^{-2} 1.0000	1.0100×10^{-2} 1.0000	1.0100×10^{-2} 1.0000
	10^{-1}	2.0050×10^{-2}	2.0100×10^{-2} 1.0025	2.0002×10^{-2} 0.9976	2.0049×10^{-2} 1.0000	2.0050×10^{-2} 1.0000
	1	1.0150	1.0774 1.0615	0.9311 0.9173	1.0046 0.9897	1.0152 1.0002
	10	1.0051×10^2	1.1105×10^2 1.1049	9.1577×10^1 0.9111	9.9945×10^1 0.9944	1.0084×10^2 1.0033

Table 2: Result of Method(I): $N = 2$, $\omega = 1$ and $\omega_0 = 10^2, 1, 10^{-2}$ for $10^{-3} \leq \lambda \leq 10$. The error is within 1.1(0.3)% under the 2(3)-loop approximation for a whole coupling region.

•Method(II): in (80), the saddle points are given by the gap equation,

$$\tilde{f}^{(1)}(t) \Big|_{t_c} = t_c - \frac{\lambda}{\omega + \lambda t_c} - \frac{1}{N} \frac{\lambda}{\omega_0 + \lambda t_c} = 0 , \quad (100)$$

yielding to

$$\Omega_c - \omega - \frac{\lambda^2}{\Omega_c} - \frac{1}{N} \frac{\lambda^2}{\Omega_c + \delta\omega} = 0 , \quad \Omega_c \equiv \omega + \lambda t_c , \quad (101)$$

with $\delta\omega$ being given by (88). This is a cubic equation of Ω_c ,

$$(\Omega_c + \delta\omega) \left((\Omega_c)^2 - \omega\Omega_c - \lambda^2 \right) = \frac{\lambda^2}{N} \Omega_c , \quad (102)$$

contrary to Method(I), where it was quadratic, (83). We write

$$\tilde{f}^{(n)}(t_c) \equiv \tilde{f}_c^{(n)} . \quad (103)$$

The stability condition is fulfilled,

$$\tilde{f}_c^{(2)} = 1 + \frac{\lambda^2}{(\Omega_c)^2} + \frac{1}{N} \frac{\lambda^2}{(\Omega_c + \delta\omega)^2} > 0 , \quad (104)$$

for any (three) saddle points Ω_c . The values of $\tilde{f}(t)$ and derivatives at Ω_c are given

$$\tilde{f}_c = \frac{1}{2\lambda^2} (\Omega_c - \omega)^2 - \ln \Omega_c - \frac{1}{N} \ln(\Omega_c + \delta\omega) , \quad (105)$$

$$\tilde{f}_c^{(n)} = (-)^n (n-1)! \left[\left(\frac{\lambda}{\Omega_c} \right)^n + \frac{1}{N} \left(\frac{\lambda}{\Omega_c + \delta\omega} \right)^n \right] ; \quad n \geq 3 . \quad (106)$$

Then from (35)

$$\begin{aligned} & \frac{\tilde{F}_{n_j+3}}{(n_j+3)!} \\ &= \frac{[-\lambda\epsilon(\Omega_c)\epsilon(\Omega_c + \delta\omega)]^{n_j+3}}{n_j+3} \frac{(\Omega_c + \delta\omega)^{n_j+3} + \frac{1}{N} (\Omega_c)^{n_j+3}}{\left[\left((\Omega_c)^2 + \lambda^2 \right) (\Omega_c + \delta\omega)^2 + \frac{\lambda^2}{N} (\Omega_c)^2 \right]^{(n_j+3)/2}} , \end{aligned} \quad (107)$$

to give, with using the condition $\sum_{j=1}^k n_j = 2L - k$,

$$\begin{aligned} I_N &\cong \sqrt{\frac{2\pi}{N}} \exp \left[-\frac{N (\Omega_c - \omega)^2}{2\lambda^2} \right] \frac{(\Omega_c)^N (\Omega_c + \delta\omega)}{\sqrt{1 + (\lambda/\Omega_c)^2 + \frac{1}{N} (\lambda/(\Omega_c + \delta\omega))^2}} \\ &\times \sum_{L=0}^{\infty} \frac{1}{N^L} \sum_{k=0}^{2L} \frac{(-)^k}{k!} (2(L+k)-1)!! \frac{\lambda^{2(L+k)}}{\left[\left((\Omega_c)^2 + \lambda^2 \right) (\Omega_c + \delta\omega)^2 + \frac{\lambda^2}{N} (\Omega_c)^2 \right]^{L+k}} \\ &\times \sum_{\substack{\sum_{j=1}^k n_j = 2L-k \\ \text{all possible } \{n_j\}}} \prod_{j=1}^k \frac{1}{n_j+3} \left[(\Omega_c + \delta\omega)^{n_j+3} + \frac{1}{N} (\Omega_c)^{n_j+3} \right] . \end{aligned} \quad (108)$$

Z , (78), is expressed, therefore, by a product of a prefactor $\mathcal{P}$ and a loop factor $\mathcal{L}$,

$$Z \equiv \mathcal{P} \times \mathcal{L} , \quad (109)$$

with

$$\mathcal{P} \equiv \exp \left[-\frac{N (\Omega_c - \omega)^2}{2\lambda^2} \right] \frac{(\Omega_c)^N (\Omega_c + \delta\omega)}{\sqrt{1 + (\lambda/\Omega_c)^2 + \frac{1}{N} (\lambda/(\Omega_c + \delta\omega))^2}} ; \quad (110)$$

$$\begin{aligned} \mathcal{L} &\equiv \sum_{L=0}^{\infty} \frac{1}{N^L} \sum_{k=0}^{2L} \frac{(-)^k}{k!} (2(L+k)-1)!! \frac{\lambda^{2(L+k)}}{\left[\left((\Omega_c)^2 + \lambda^2 \right) (\Omega_c + \delta\omega)^2 + \frac{\lambda^2}{N} (\Omega_c)^2 \right]^{L+k}} \\ &\times \sum_{\substack{\sum_{j=1}^k n_j = 2L-k \\ \text{all possible } \{n_j\}}} \prod_{j=1}^k \frac{1}{n_j+3} \left[(\Omega_c + \delta\omega)^{n_j+3} + \frac{1}{N} (\Omega_c)^{n_j+3} \right] . \end{aligned} \quad (111)$$

Now solve the gap equation to find that there are three kinds of $1/N$ series,

$$\Omega_c^{(i)} = \Omega_0^{(i)} + \sum_{j=1} \frac{\Omega_j^{(i)}}{N^j} ; \quad (i = 1, 2, 3) . \quad (112)$$

where

$$\Omega_0^{(i)} = \begin{cases} \Omega_0^{(\pm)} ; & i = 1, 2 \\ -\delta\omega ; & i = 3 \end{cases} , \quad (113)$$

with $\Omega_0^{(\pm)}$ being given by (84), then up to $O(1/N^2)$,

$$\Omega_1^{(i)} = \frac{\lambda^2 \Omega_0^{(i)}}{B_i} ; \quad \Omega_2^{(i)} = \frac{\lambda^2 \Omega_1^{(i)} - \left(\Omega_1^{(i)}\right)^2 A_i}{B_i} . \quad (114)$$

$$A_i \equiv 3\Omega_0^{(i)} - \omega + \delta\omega ; \quad B_i \equiv 3\left(\Omega_0^{(i)}\right)^2 - 2(\omega - \delta\omega)\Omega_0^{(i)} - \omega\delta\omega - \lambda^2 . \quad (115)$$

These are sufficient under the 3-loop approximation ($O(1/N^2)$). Accordingly write Z as $Z^{(i)}$ in (109) such that

$$Z^{(i)} = \mathcal{P}^{(i)} \times \mathcal{L}^{(i)} ; \quad (i = 1, 2, 3) , \quad (116)$$

with the prefactor,

$$\mathcal{P}^{(i)} \equiv F^{(i)}(1)F^{(i)}(2)F^{(i)}(3) \left(\Omega_c^{(i)} + \delta\omega\right) , \quad (117)$$

where

$$F^{(i)}(1) \equiv \exp \left[-N \frac{\left(\Omega_c^{(i)} - \omega\right)^2}{2\lambda^2} \right] ; \quad F^{(i)}(2) \equiv \left(\Omega_c^{(i)}\right)^N , \quad (118)$$

$$F^{(i)}(3) \equiv \left[1 + \left(\frac{\lambda}{\Omega_c^{(i)}}\right)^2 + \frac{1}{N} \left(\frac{\lambda}{\Omega_c^{(i)} + \delta\omega}\right)^2 \right]^{-1/2} , \quad (119)$$

and the loop factor, $\mathcal{L}^{(i)}$, up to $O(1/N^2)$,

$$\begin{aligned} \mathcal{L}^{(i)} &\equiv \sum_{L=0}^{\infty} \frac{1}{N^L} \sum_{k=0}^{2L} \frac{(-)^k}{k!} (2(L+k)-1)!! \left(\frac{\lambda^2}{F^{(i)}(4)}\right)^{L+k} \sum_{\text{all possible } \{n_j\}}^{\sum_{j=1}^k n_j = 2L-k} \prod_{j=1}^k \frac{F^{(n_j+3;i)}(5)}{n_j+3} \\ &= 1 + \frac{\lambda^2}{N F^{(i)}(4)} \left[-\frac{3}{4} \frac{\lambda^2 F^{(4;i)}(5)}{F^{(i)}(4)} + \frac{5}{6} \left(\frac{\lambda^2 F^{(3;i)}(5)}{F^{(i)}(4)} \right)^2 \right] \\ &\quad + \frac{\lambda^4}{(N F^{(i)}(4))^2} \left[-\frac{5}{2} \frac{\lambda^2 F^{(6;i)}(5)}{F^{(i)}(4)} + 7 \frac{\lambda^4 F^{(3;i)}(5) F^{(5;i)}(5)}{(F^{(i)}(4))^2} + \frac{105}{32} \left(\frac{\lambda^2 F^{(4;i)}(5)}{F^{(i)}(4)} \right)^2 \right. \\ &\quad \left. - \frac{105}{8} \frac{\left(F^{(3;i)}(5)\right)^2 F^{(4;i)}(5)}{(F^{(i)}(4))^3} + \frac{385}{72} \left(\frac{\lambda^2 F^{(3;i)}(5)}{F^{(i)}(4)} \right)^4 \right] , \end{aligned} \quad (120)$$

where

$$F^{(i)}(4) \equiv \left(\left(\Omega_c^{(i)} \right)^2 + \lambda^2 \right) \left(\Omega_c^{(i)} + \delta\omega \right)^2 + \frac{\lambda^2}{N} \left(\Omega_c^{(i)} \right)^2 ; \quad (121)$$

$$F^{(M;i)}(5) \equiv \left(\Omega_c^{(i)} + \delta\omega \right)^M + \frac{1}{N} \left(\Omega_c^{(i)} \right)^M ; \quad M = 3, 4, \dots, 6 . \quad (122)$$

Note that

$$\Omega_c^{(i)} + \delta\omega = \begin{cases} \left(\Omega_0^{(\hat{i})} + \delta\omega \right) \left[1 + \frac{1}{N} \frac{\Omega_1^{(\hat{i})}}{\Omega_0^{(\hat{i})} + \delta\omega} + \frac{1}{N^2} \frac{\Omega_2^{(\hat{i})}}{\Omega_0^{(\hat{i})} + \delta\omega} \right] ; & \hat{i} = 1, 2 \\ \frac{\Omega_1^{(3)}}{N} \left[1 + \frac{1}{N} \frac{\Omega_2^{(3)}}{\Omega_1^{(3)}} \right] ; & i = 3 \end{cases} , \quad (123)$$

so that the contribution of the third saddle, $i = 3$, starts from $O(1/N)$, that is, 2-loop. Therefore we write

$$\mathcal{P}^{(i)} = \begin{cases} \mathcal{P}_0^{(\hat{i})} \left[1 + \frac{\mathcal{P}_1^{(\hat{i})}}{N} + \frac{\mathcal{P}_2^{(\hat{i})}}{N^2} \right] ; & \hat{i} = 1, 2 \\ \frac{\mathcal{P}_1^{(3)}}{N} \left[1 + \frac{\mathcal{P}_2^{(3)}}{N} \right] ; & i = 3 \end{cases} ; \quad (124)$$

and

$$\mathcal{L}^{(i)} = \begin{cases} \mathcal{L}_0^{(\hat{i})} + \frac{\mathcal{L}_1^{(\hat{i})}}{N} + \frac{\mathcal{L}_2^{(\hat{i})}}{N^2} ; & \hat{i} = 1, 2 \\ \mathcal{L}_0^{(3)} + \frac{\mathcal{L}_1^{(3)}}{N} ; & i = 3 \end{cases} . \quad (125)$$

Explicit forms of those functions, $\mathcal{P}_0^{(\hat{i})} \sim \mathcal{P}_1^{(3)}$ and $\mathcal{L}_0^{(\hat{i})} \sim \mathcal{L}_1^{(3)}$, are (after lengthy calculation) given in the appendix C.

The tree part is, from (169) and (172),

$$\begin{aligned} Z_{\text{tree}}^{(\hat{i})} &= F_0^{(\hat{i})}(1) F_0^{(\hat{i})}(2) \left(\Omega_0^{(\hat{i})} + \delta\omega \right) \Big|_{\Omega_1^{(\hat{i})} \mapsto 0} \\ &= \exp \left[-\frac{N}{2\lambda^2} \left(\Omega_0^{(\hat{i})} - \omega \right)^2 \right] \left(\Omega_0^{(\hat{i})} \right)^N \left(\Omega_0^{(\hat{i})} + \delta\omega \right) . \end{aligned} \quad (126)$$

Since $Z_{\text{tree}}^{(3)} = 0$, this is equivalent to the one, (94), in Method(I). The 1-loop part reads

$$Z_{1\text{-loop}}^{(\hat{i})} = \mathcal{P}_0^{(\hat{i})} \mathcal{L}_0^{(\hat{i})} \Big|_{\Omega_1^{(\hat{i})} \mapsto 0} = \epsilon \left(\Omega_0^{(\hat{i})} \right) \exp \left[-\frac{N \left(\Omega_0^{(\hat{i})} - \omega \right)^2}{2\lambda^2} \right] \frac{\left(\Omega_0^{(\hat{i})} \right)^{N+1} \left(\Omega_0^{(\hat{i})} + \delta\omega \right)}{\sqrt{\left(\Omega_0^{(\hat{i})} \right)^2 + \lambda^2}} , \quad (127)$$

and $Z_{1\text{-loop}}^{(3)} = 0$ with the aid of (188) and (198), which again matches with (96). Next

$$Z_{2\text{-loop}}^{(\hat{i})} = \mathcal{P}_0^{(\hat{i})} \left[1 + \frac{\left(\mathcal{P}_1^{(\hat{i})} + \mathcal{L}_1^{(\hat{i})} \right)}{N} \right] \Big|_{\Omega_2^{(\hat{i})} \mapsto 0} = Z_{1\text{-loop}}^{(\hat{i})} \left[1 + \frac{\left(\mathcal{P}_1^{(\hat{i})} + \mathcal{L}_1^{(\hat{i})} \right)}{N} \right] \Big|_{\Omega_2^{(\hat{i})} \mapsto 0} ; \quad (128)$$

where

$$\begin{aligned}
\left(\mathcal{P}_1^{(\hat{i})} + \mathcal{L}_1^{(\hat{i})}\right) \Big|_{\Omega_2^{(\hat{i})} \mapsto 0} &= \left(\sum_{r=1}^3 F_1^{(i)}(r) + \frac{\Omega_1^{(\hat{i})}}{\Omega_0^{(\hat{i})} + \delta\omega} + \mathcal{L}_1^{(\hat{i})}\right) \Big|_{\Omega_2^{(\hat{i})} \mapsto 0} \\
&= -\frac{\left(\Omega_1^{(\hat{i})}\right)^2}{2\lambda^2} - \frac{1}{2} \left(\frac{\Omega_1^{(\hat{i})}}{\Omega_0^{(\hat{i})}}\right)^2 - \frac{\lambda^2}{\left(\Omega_0^{(\hat{i})}\right)^2 + \lambda^2} \left[\frac{\Omega_1^{(\hat{i})}}{\Omega_0^{(\hat{i})}} - \frac{1}{2} \left(\frac{\Omega_0^{(\hat{i})}}{\Omega_0^{(\hat{i})} + \delta\omega}\right)^2\right] \\
&\quad + \frac{\Omega_1^{(\hat{i})}}{\Omega_0^{(\hat{i})} + \delta\omega} - \frac{3\lambda^4}{4 \left[\left(\Omega_0^{(\hat{i})}\right)^2 + \lambda^2\right]^2} + \frac{5\lambda^6}{6 \left[\left(\Omega_0^{(\hat{i})}\right)^2 + \lambda^2\right]^3}, \tag{129}
\end{aligned}$$

from (170), (172), (176) and (199). For $i = 3$ we obtain

$$\begin{aligned}
Z_{2\text{-loop}}^{(3)} &= \mathcal{P}_1^{(3)} \mathcal{L}_0^{(3)} \Big|_{2\text{-loop}} \\
&= \epsilon \left(\Omega_1^{(3)}\right) \exp \left[-\frac{N \left(\Omega_0^{(3)} - \omega\right)^2}{2\lambda^2} \right] \frac{\left(\Omega_0^{(3)}\right)^N \left(\Omega_1^{(3)}\right)^2}{e} \frac{1}{\sqrt{N}\lambda} \frac{1}{N} \frac{13}{12}, \tag{130}
\end{aligned}$$

in view of (189) and (201).

Finally Z under 3-loop for $i = 1, 2$ is

$$\begin{aligned}
Z_{3\text{-loop}}^{(\hat{i})} &= \mathcal{P}_0^{(\hat{i})} \left[1 + \frac{\mathcal{P}_1^{(\hat{i})} + \mathcal{L}_1^{(\hat{i})}}{N} + \frac{\mathcal{P}_2^{(\hat{i})} + \mathcal{L}_2^{(\hat{i})} + \mathcal{P}_1^{(\hat{i})} \mathcal{L}_1^{(\hat{i})}}{N^2} \right] \\
&= Z_{1\text{-loop}}^{(\hat{i})} \left[1 + \frac{\mathcal{P}_1^{(\hat{i})} + \mathcal{L}_1^{(\hat{i})}}{N} + \frac{\mathcal{P}_2^{(\hat{i})} + \mathcal{L}_2^{(\hat{i})} + \mathcal{P}_1^{(\hat{i})} \mathcal{L}_1^{(\hat{i})}}{N^2} \right], \tag{131}
\end{aligned}$$

whose functions $\mathcal{P}_{1,2}^{(\hat{i})}, \mathcal{L}_{1,2}^{(\hat{i})}$ are given in the appendix C; (191), (193), (199), and (200).

Meanwhile $i = 3$ is

$$\begin{aligned}
Z_{3\text{-loop}}^{(3)} &= \frac{\mathcal{P}_1^{(3)}}{N} \left(1 + \frac{\mathcal{P}_2^{(3)}}{N} \right) \left(\mathcal{L}_0^{(3)} + \frac{\mathcal{L}_1^{(3)}}{N} \right) \Big|_{3\text{-loop}} = \frac{\mathcal{P}_1^{(3)}}{N} \left[\mathcal{L}_0^{(3)} + \frac{\mathcal{L}_1^{(3)} + \mathcal{L}_0^{(3)} \mathcal{P}_2^{(3)}}{N} \right] \Big|_{3\text{-loop}} \\
&= \epsilon \left(\Omega_1^{(3)}\right) \exp \left[-\frac{N \left(\Omega_0^{(3)} - \omega\right)^2}{2\lambda^2} \right] \frac{\left(\Omega_0^{(3)}\right)^N \left(\Omega_1^{(3)}\right)^2}{e\sqrt{N}\lambda} \\
&\quad \times \left[\frac{1}{N} \frac{313}{288} + \frac{1}{N^2} \left(-\frac{13}{12} \left[\left(\frac{\Omega_1^{(3)}}{\Omega_0^{(3)}}\right)^2 + \frac{\left(\Omega_1^{(3)}\right)^2}{\lambda^2} \right] + \frac{313}{288} \mathcal{P}_2^{(3)} \right) \right], \tag{132}
\end{aligned}$$

where $\mathcal{P}_2^{(3)}$ is given in (192) and use has been made of (203) and (204). For Z_{tree} and $Z_{1\text{-loop}}$, there is no difference from Method (I). However, in 2- and 3-loop, the third saddle starts contributing to give

$$Z_{l\text{-loop}}^{\text{Total}} = Z_{l\text{-loop}}^{(1)} + Z_{l\text{-loop}}^{(2)} + Z_{l\text{-loop}}^{(3)}; \quad (l = 2, 3), \tag{133}$$

whose numerical results, when $N = 2$ and $\omega = 1$ with $\omega_0 = 10^2$ and 1, are equivalent to those of Method(I) for $10^{-3} \leq \lambda \leq 10$, which is in the table 2. In the table 3, we list the result of $\omega_0 = 10^{-2}$, in which we see disparities at 2- and 3-loop in the weak coupling region $10^{-2} \leq \lambda \leq 1$. Discrepancies are notable, reaching to ~ 600 times to the exact value at $\lambda = 10^{-1}$ in 3-loop.

$\omega_0 = 10^{-2}$	λ	Exact	tree (tree)/Exact	1-loop (1-loop)/Exact	2-loop (2-loop)/Exact	3-loop (3-loop)/Exact
	10^{-3}	1.0001×10^{-2}	1.0001×10^{-2} 1.0000	1.0001×10^{-2} 1.0000	1.0001×10^{-2} 1.0000	1.0001×10^{-2} 1.0000
	10^{-2}	1.0101×10^{-2}	1.0101×10^{-2} 1.0000	1.0100×10^{-2} 1.0000	9.6026×10^{-3} 0.9507	1.0335×10^{-2} 1.0232
	10^{-1}	2.0050×10^{-2}	2.0100×10^{-2} 1.0025	2.0002×10^{-2} 0.9976	-3.1834×10^{-1} -15.877	1.2398×10^1 618.36
	1	1.0150	1.0774 1.0615	0.9311 0.9173	0.8719 0.8590	1.2693 1.2506
	10	1.0051×10^2	1.1105×10^2 1.1049	9.1577×10^1 0.9111	9.9931×10^1 0.9942	1.0085×10^2 1.0034

Table 3: Result of Method(II): $N = 2$, $\omega = 1$ and $\omega_0 = 10^{-2}$ for $10^{-3} \leq \lambda \leq 10$. Discrepancies to Method(I), in the table 2, are acknowledged in 2, 3-loop for $10^{-2} \leq \lambda \leq 10$.

The reason can be seen from $Z_{3\text{-loop}}^{(3)}$ (130) by putting $\omega_0 \ll 1$;

$$Z_{3\text{-loop}}^{(3)} \sim \exp \left[-N \frac{\omega_0^2}{2\lambda^2} \right] \frac{1}{\lambda} ; \quad (134)$$

which has some peak around $\lambda \sim \omega_0$. The graph is shown in the figure 2, implying a large deviation at the value $\omega_0 = 0.01$. Although the case would not be included in the real




Figure 2: The graph of $Z_{3\text{-loop}}^{(3)}$ is shown with $N = 2, \omega = 1$ for $0 \leq \lambda \leq 1$. The vertical line is the value of $Z_{3\text{-loop}}^{(3)}$ itself. The solid, dotted, and dashed line designate $\omega_0 = 0.01, 0.05, 0.1$. The deviation from the exact value (which is almost zero) is seen around $\lambda \sim 1.5$ when $\omega_0 = 0.01$.

physical situation, u- and d-quarks are lighter than s-quark, Method(II) is worse than (I) in the above situation. Therefore, a recipe for an approximation Method(I) is better than (II) and moreover simpler for actual calculations.

5 Discussion

Under Method(II), we encounter the same situation in the bosonic four-body model under the weak coupling region when the "mass", ω_0 , is tiny: consider σ (8) in the introduction as bosonic variables, to have

$$Z^\omega \sim \int \frac{dy}{\omega_0 + i\lambda y} \exp[-Nf(y)] , \quad (135)$$

where in view of (4) and (6),

$$f(y) = \frac{y^2}{2} + \ln(\omega_0 + i\lambda y) . \quad (136)$$

(We have omitted irrelevant factors, $2\pi, N$.) Therefore

$$Z^\omega \sim \int dt \exp[-N\tilde{f}(t)] , \quad (137)$$

$$\tilde{f}(t) \equiv f(t) + \frac{1}{N} \ln g(t) ; \quad g(t) \equiv \omega_0 + i\lambda t . \quad (138)$$

(Again we have switched, $y \mapsto t$.)

Take, for the time being, $\tilde{f}(t), f(t), g(t)$ as generic, in other words, start from (14) to make a general discussion: there emerge additional saddle points, t_0^A , in the gap equation when $N \mapsto \infty$,

$$0 = f'(t) + \frac{1}{N} \frac{g'(t)}{g(t)} \xrightarrow{N \mapsto \infty} g(t)f'(t) = 0 ; \quad (139)$$

other than $f'(t_0) = 0$ such that

$$g(t_0^A) = 0 . \quad (140)$$

Therefore the additional saddle point(s) is expanded as

$$t_c = t_0^A + \frac{t_1^A}{N} + O\left(\frac{1}{N^2}\right) , \quad (141)$$

around which

$$g_c \equiv g(t_c) = \frac{t_1^A}{N} g_A^{(1)} + O\left(\frac{1}{N^2}\right) ; \quad g_A^{(1)} \equiv g^{(1)}(t_0^A) . \quad (142)$$

Now recall that one of the 2-loop terms is given as ((35) with (36))

$$Z_{2\text{-loop}}^\omega \sim \frac{1}{\sqrt{N\tilde{f}_c^{(2)}}} \exp[-N\tilde{f}_c] \frac{\tilde{f}_c^{(4)}}{N\tilde{f}_c^{(2)}} = \frac{\tilde{f}_c^{(4)}}{g_c (N\tilde{f}_c^{(2)})^{3/2}} \exp[-Nf_c] . \quad (143)$$

By noting

$$\tilde{f}_c^{(n)} = f_c^{(n)} + \frac{1}{N} \left[(-)^{n-1} n! \left(\frac{g_c^{(1)}}{g_c} \right)^n + O\left(\frac{1}{(g_c)^{n-1}}\right) \right] , \quad (144)$$

we find

$$Z_{2\text{-loop}}^\omega \sim \frac{\tilde{f}_c^{(4)}}{g_c \left(N \tilde{f}_c^{(2)} \right)^{3/2}} \exp[-N f_c] \sim \frac{g_c^{(1)}}{(g_c)^2} \exp[-N f_c] , \quad (145)$$

around the additional saddle point, which reads in the bosonic case (138)

$$Z_{2\text{-loop}}^\omega \sim \frac{g_c^{(1)}}{(g_c)^2} \exp[-N f_c] \stackrel{\omega_0 \rightarrow 0}{\sim} \frac{1}{\lambda} , \quad (146)$$

since $g_c \sim g_c^{(1)} \sim \lambda$ (138), again implying a large deviation.

In summary, Method(II) was superficially simpler than (I) but needs a rather cumbersome procedure in the actual calculation and moreover always seems to suffer from a large deviation when in a weak coupling region when the one "mass" ω_0 is tiny.

As the final comment, we check the validity of Method(I) in an alternative way: the case of $\omega_0 = \omega$ corresponds to the $N = 3$ version of the model[8],

$$Z_{\text{KS}} \equiv \int d^N \hat{\xi} d^N \hat{\xi}^* \exp \left[-\hat{\xi}^* \cdot \omega \cdot \hat{\xi} + \frac{(\lambda_{\text{KS}})^2}{2N} \left(\hat{\xi}^* \cdot \hat{\xi} \right)^2 \right] . \quad (147)$$

So if we put

$$(\lambda_{\text{KS}})^2 = \frac{3}{2} \lambda^2 , \quad (148)$$

this should agree with our model($N = 2$). Applying a usual AFM (that is, from the expression (1) to (7)), we have results up to 3-loop, which is listed in the table 4. By comparing this with the table 2 (of $\omega_0 = \omega$), there is no big difference: almost all data show that the standard treatment of the table 4 yields a slightly better value except the tree approximation in $10^{-1} \leq \lambda \leq 10$ where our model under Method(I) results better.

λ	Exact	tree (tree)/Exact	1-loop (1-loop)/Exact	2-loop (2-loop)/Exact	3-loop (3-loop)/Exact
$\sqrt{\frac{3}{2}} \times 10^{-3}$	1.0000	1.0000 1.0000	1.0000 1.0000	1.0000 1.0000	1.0000 1.0000
$\sqrt{\frac{3}{2}} \times 10^{-2}$	1.0002	1.0002 1.0001	1.0002 1.0000	1.0002 1.0000	1.0002 1.0000
$\sqrt{\frac{3}{2}} \times 10^{-1}$	1.0150	1.0224 1.0073	1.0151 1.0001	1.0150 1.0000	1.0150 1.0000
$\sqrt{\frac{3}{2}}$	2.5000	3.0574 1.2230	2.5433 1.0173	2.5032 1.0013	2.4996 0.9998
$\sqrt{\frac{3}{2}} \times 10$	1.5100×10^2	2.0188×10^2 1.3369	1.5484×10^2 1.0254	1.5106×10^2 1.0004	1.5089×10^2 0.9993

Table 4: Results of the model (147) at $N = 3$, $\omega = 1$, and $\lambda_{\text{KS}} = \sqrt{\frac{3}{2}} \lambda$.

Acknowledgements

The authors are grateful to H. So for discussions, especially for guiding them to the second proof of (54).

A The table of $T(L, k|2L - k)$ and $T(L, k|2L - k - 1)$ defined by eq.(52)

In this appendix, we list the table of $T(L, k|2L - k) / (2(L + k) - 1)!!$ instead of $T(L, k|2L - k)$ itself; since otherwise the values become very large. The range up to $L = 14$ is needed for the calculation of the Gamma function in Sec.3 and to $L = 4$ in Sec.4.

$k \setminus L$	0	1	2	3	4	5	6	7
0	1	0	0	0	0	0	0	0
1		-1/4	-1/6	-1/8	-1/10	-1/12	-1/14	-1/16
2		1/18	47/480	153/1400	3349/30240	42131/388080	605453/5765760	655217/6486480
3			-1/72	-493/17280	-4049/100800	-59197/1209600	-161453/2910600	-81158813/1345344000
4			1/1944	77/25920	190261/29030400	1722811/163296000	406957909/27941760000	668430857/36324288000
5				-1/7776	-503/933120	-143921/116121600	-5689699/2612736000	-39726066467/12070840320000
6				1/524880	107/4665600	44021/522547200	74863031/376233984000	4489161401/12070840320000
7					-1/2099520	-1699/503884800	-216793/18811699200	-13916027/501645312000
8					1/264539520	137/1763596800	282133/658409472000	142181689/101583175680000
9						-1/1058158080	-643/63489484800	-1143257/23702740992000
10						1/214277011200	167/1142810726400	361657/319987003392000
11							-1/857108044800	-29/1645647446016
12							1/254561089305600	197/1131382619136000
13								-1/101824435722400
14								1/416971064282572800

$k \setminus L$	8	9	10
0	0	0	0
1	-1/18	-1/20	-1/22
2	23763863/245044800	158899519/1707145440	11098301/124156032
3	-232229821/3632428800	-1974182737/29640619008	-176989210093/2581203905280
4	3218616617/146459529216	131020195003/5187108326400	1429205465892419/50591596543488000
5	-59153229587/13076743680000	-237186996829/40683202560000	-4136077769339/576345369600000
6	449840404627/747242496000000	2187240114496949/247150455520000000	27225846275010179/22408307970048000000
7	-15727453241/289700167680000	-11643095464009/125536739328000000	-711993190614367/4943009111040000000
8	570158851513/166867296583680000	23485635969143/3389491961856000000	4800125589203449/388250897448960000000
9	-434386633/2844328919040000	-754670748947/2002407559004160000	-28426615868773/36154580926464000000
10	1125369389/230390642442240000	4768426956641/315379190543155200000	61720076581079809/1639971790824407040000000
11	-142837/1279948013568000	-414828299/921562569768960000	-46647858451427/34060952578660761600000
12	2278697/1267148533432320000	64781047/6516763886223360000	14267906296747/374670478365268377600000
13	-347/17455617552384000	-7370813/45617347203563520000	-147673007/182469388814254080000
14	227/1588461197266944000	563401/296512756823162880000	8377793029/640467554738031820800000
15	-1/1667884257130291200	-6271/400292221711269888000	-1697149/10674459245633863680000
16	1/900657498850357248000	257/3002191662834524160000	957911/672490932474933411840000
17		-1/3602629995401428992000	-983/108078899862042869760000
18		1/2480410751833883860992000	41/1049909312945559306240000
19			-1/9921643007335535443968000
20			1/8483004771271882804592640000

Table 5: The values of $T(L, k|2L - k) / (2(L + k) - 1)!!$ for $L = 0 \sim 10$ in sec.3 as well as sec.4

B The proof of the relation (49) and of (54)

In this appendix, we first prove (49):

$$\sum_{k=0}^{2L} T(L, k | \leq 2L - k) = \sum_{k=0}^{2L} T(L, k | 2L - k) , \quad (149)$$

where

$$T(L, k | \leq 2L - k) = (-)^k (2(L + k) - 1)!! \sum_{\substack{\text{all possible } \{A_\alpha\} \\ \sum_{\alpha=1}^P Q_\alpha = k ; (P \leq k)}} \frac{\sum_{\alpha=1}^P Q_\alpha A_\alpha \leq 2L - k}{Q_1! \cdots Q_P!} \times \frac{1}{(A_1 + 3)^{Q_1} (A_2 + 3)^{Q_2} \cdots (A_P + 3)^{Q_P}} , \quad (150)$$

$k \setminus L$	11	12
0	0	0
1	-1/24	-1/26
2	265842403/3093554464	20666950267/249864014400
3	-4160074481/59435616240	-205489485020713/2894174878795200
4	2665668285176389/86090742017280000	15599534488926830887/46677252445208956000
5	-8637934672612781/1011831930869760000	-40541391139263449/4097919320022528000
6	203058721852862831/128059978750704000000	40047282902251643797/20117058480110592000000
7	-841208011558860343/4033495434608640000000	-93845936786285052059/327833545601802240000000
8	405347221415512403/20167477173043200000000	3993321076456790080543/131133418240720896000000000
9	-198721338986302927/136664315902033920000000	-3058025156141685767/1244621448393523200000000
10	691811729911733023/8609851901828136960000000	21345802689252486829/139397602220074598400000000
11	-202677271543968731/59038984469678653440000000	-1543928961441550217/206636445643875287040000000
12	3374227932094529/29519492234839326720000000	19789232975365781149/68666880398580095385600000000
13	-40177324205483/13488137221149661593600000	-103383349078912121/11689718924996373381120000000
14	79944551761007/1315093379062092005376000000	296393081576585261/1367697114224575685591040000000
15	-2481097459/2561870218952127283200000	-22322956680727/5260373516248368021504000000
16	2893597613/242096735690976028262400000	14641217114597/220935687682431456903168000000
17	-101081/896654576633244549120000	-794890043/968386942763904113049600000
18	74363/93537375153331647283200000	5917683217/740816011214386646482944000000
19	-9637/2381194321760528506552320000	-6685391/111122401682157996972441600000
20	317/22621346056725020812247040000	56741/165594167212627603331481600000
21	-1/33932019085087531218370560000	-61/42861497791689513117941760000
22	1/3527233838948488701496197120000	347/85508688094420578670293811200000
23		-1/141089335355793954805984788480000
24		1/175232954511896091869033107292160000

$k \setminus L$	13	14
0	0	0
1	-1/28	-1/30
2	192066102203/2409402996000	5733412167187/74530866009600
3	-316144708749043/4410171243878400	-5230982263631/72535711248000
4	1039787017585599191/29173282778255616000	86993393978615452781/2310523996037844787200
5	-1321164558963178417729/117626676161926643712000	-409853074642336437191/32674076711646289920000
6	124759585247032919932373/51461670820842906624000000	515724686858368491454901/178997115898584023040000000
7	-3511001126573317919533/9317374453945958400000000	-26661050735382530348701583/55578604486510339153920000000
8	9940709833698231428297/227610147374965555200000000	15680328374701897384850153/261546374054166301900800000000
9	-41120299919631976751017/10621806877498392576000000000	-91936525080313650485417/15932710316247588864000000000
10	136358632264218467927423/50984673011992284364800000000	12636723159765370307618303/28980761501553509007360000000000
11	-1026261079863684476899/70256391518917597593600000000	-40157529586233540547571/1529540190359768530944000000000
12	2015526363139860379999/3161537618351291891712000000000	23571578581358615769453103/18501318142591760150298624000000000
13	-880105003537523330687/39277455587987814560563200000000	-83901448111763933230733/16692918624894821188239360000000000
14	338122116167654792537/530245650437835496567603200000000	884333215764851072621/5450748938741166102282240000000000
15	-26876946960034477/1823596152299434247454720000000	-1186820857702705430201/27572773822767445821515366400000000
16	1032930932083247/3730083038794297324339200000000	1167709777320002702963/1240774822024535061968191488000000000
17	-302359614816397/715831628091077920366264320000000	-526519789986178363/310193705506133765492047872000000000
18	31701146999281/608456883877416232311324672000000	266111014317840767/1054658598720854802672962764800000000
19	-4588191221/8889792134572639757795328000000	-40513735354423/13142668691752190617924612915200000
20	9262374577/2280231682517882097874501632000000	576706185137543/1872830288574687163054257340416000000
21	-7598699/304030891002384279716600217600000	-2275833187/9120926730071528391498006528000000
22	5619821/47884865332875524055364534272000000	1596747533/985060086847725066281784705024000000
23	-13723/33861440485390549153436349235200000	-17489411/2106934074646523058436039507968000000
24	29/29954351198614716558809078169600000	14256551/436135353451830273096260178149376000000
25	-1/700931818047584367476132429168640000	-211/2213468899097634844661470828953600000
26	1/1025112783894592137433843677659136000000	37/191163223103886645675308844318720000000
27		-1/4100451135578368549735374710636544000000
28		1/6974867381618804903099872382792761344000000

Table 6: The values of $T(L, k|2L - k)/(2(L + k) - 1)!!$ for $L = 11 \sim 14$ in sec.3 as well as sec.4

and $T(L, k | 2L - k)$ in RHS is given in terms of the conditional sum $\sum_{\alpha=1}^P Q_{\alpha} A_{\alpha} = 2L - k$ instead of $\sum_{\alpha=1}^P Q_{\alpha} A_{\alpha} \leq 2L - k$. When $L = 0$ (which also means $k = 0$) the relation (149) holds

$$T(0, 0 | \leq 0) = T(0, 0 | 0) , \quad (151)$$

since $\sum_{\alpha=1}^P Q_{\alpha} A_{\alpha} \leq 0$ is nothing but $\sum_{\alpha=1}^P Q_{\alpha} A_{\alpha} = 0$ for any positive Q_{α}, A_{α} . Therefore we can set $L \neq 0$ in the following. Likewise when $k = 2L$

$$T(L, 2L | \leq 0) = T(L, 2L | 0) . \quad (152)$$

Therefore the target relation (149) turns out to be

$$\sum_{k=0}^{2L-1} T(L, k | < 2L - k) = 0 , \quad L \neq 0 , \quad (153)$$

with $T(L, k \mid < 2L - k)$ being given by the conditional sum $\sum_{\alpha=1}^P Q_{\alpha} A_{\alpha} < 2L - k$ in (150), which is further rewritten as

$$\begin{aligned} \text{LHS of (153)} &= T(L, 0 \mid < 2L) + T(L, 1 \mid 2L - 2) \\ &+ \sum_{k=1}^{2L-2} \left\{ T(L, k \mid \leq 2L - k - 2) + T(L, k + 1 \mid 2L - k - 2) \right\}, \end{aligned} \quad (154)$$

since with the aid of (an obvious relation)

$$T(L, k \mid < 2L - k) = T(L, k \mid 2L - k - 1) + T(L, k \mid \leq 2L - k - 2), \quad (155)$$

LHS reads

$$\begin{aligned} \text{LHS of (153)} &= T(L, 0 \mid < 2L) + \sum_{k=1}^{2L-1} \left\{ T(L, k \mid 2L - k - 1) + T(L, k \mid \leq 2L - k - 2) \right\} \\ &= T(L, 0 \mid < 2L) + T(L, 1 \mid 2L - 2) + \sum_{k=2}^{2L-1} T(L, k \mid 2L - k - 1) \\ &\quad + \sum_{k=1}^{2L-1} T(L, k \mid \leq 2L - k - 2), \end{aligned} \quad (156)$$

whose last term becomes $\sum_{k=2}^{2L-2} T(L, k \mid \leq 2L - k - 2)$, because $T(L, 2L - 1 \mid \leq -1) = 0$, giving the third term of (154). Meanwhile the third term in (156) gives the last term of (154) by shifting $k \mapsto k + 1$.

The first term cancels the second in RHS of (154); since from (150)

$$T(L, 0 \mid < 2L) = (2L - 1)!! , \quad (157)$$

and

$$T(L, 1 \mid 2L - 2) = -(2L + 1)!! \frac{1}{(2L - 2) + 3} = -(2L - 1)!! , \quad (158)$$

from (150) (with changing the conditional sum to $Q_1 A_1 = 2L - 2$) and $Q_1 = 1$ (obtaining from $\sum_{\alpha=1}^P Q_{\alpha} = 1$; $P \leq 1$). Therefore if we show

$$T(L, K + 1 \mid 2L - K - 2) = -T(L, K \mid \leq 2L - K - 2) ; \quad 1 \leq K \leq 2L - 2. \quad (159)$$

we find that RHS of (154) vanishes, accomplishing the proof of (149).

Now prove (159): LHS reads

$$\begin{aligned} T(L, K + 1 \mid 2L - K - 2) &= (-)^{K+1} (2(L + K) + 1)!! \\ &\times \sum_{\substack{\sum_{\alpha=1}^P Q_{\alpha} A_{\alpha} = 2L - K - 2 \\ \text{all possible } \{A_{\alpha}\} \\ \sum_{\alpha=1}^P Q_{\alpha} = K + 1 ; (P \leq K + 1)}} \frac{1}{Q_1! \cdots Q_P!} \frac{1}{(A_1 + 3)^{Q_1} \cdots (A_P + 3)^{Q_P}}. \end{aligned} \quad (160)$$

In view of (150), the conditional sum of RHS, $\sum_{\alpha=1}^P Q_{\alpha} A_{\alpha} \leq 2L - K - 2$, is fulfilled by putting some Q_{β} ; ($\beta \in \alpha$) to $Q_{\beta} - 1$, which brings the sum $\sum_{\alpha=1}^P Q_{\alpha} = K$ to

$\sum_{\alpha=1}^P Q_{\alpha} - 1 = K$, while keeping $\sum_{\alpha=1}^P Q_{\alpha} A_{\alpha} = 2L - K - 2$, to give

$$T(L, K \mid \leq 2L - K - 2) = (-)^K (2(L + K) - 1)!! \sum_{\substack{\sum_{\alpha=1}^P Q_{\alpha} A_{\alpha} = 2L - K - 2 \\ \text{all possible } \{A_{\alpha}\} \\ \sum_{\alpha=1}^P Q_{\alpha} = K + 1 ; (P \leq K + 1)}} \frac{1}{Q_1! \cdots (Q_{\beta} - 1)! \cdots Q_P!} \frac{1}{(A_1 + 3)^{Q_1} \cdots (A_{\beta} + 3)^{Q_{\beta-1}} \cdots (A_P + 3)^{Q_P}} . \quad (161)$$

RHS of (161) is further rewritten as

$$\begin{aligned} \text{RHS of (161)} &= (-)^K (2(L + K) - 1)!! \sum_{\substack{\sum_{\alpha=1}^P Q_{\alpha} A_{\alpha} = 2L - K - 2 \\ \text{all possible } \{A_{\alpha}\} \\ \sum_{\alpha=1}^P Q_{\alpha} = K + 1 ; (P \leq K + 1)}} \left[\sum_{\beta=1}^P Q_{\beta} (A_{\beta} + 3) \right] \\ &\quad \times \frac{1}{Q_1! \cdots Q_P!} \frac{1}{(A_1 + 3)^{Q_1} \cdots (A_P + 3)^{Q_P}} \\ &= (-)^K (2(L + K) + 1)!! \\ &\quad \times \sum_{\substack{\sum_{\alpha=1}^P Q_{\alpha} A_{\alpha} = 2L - K - 2 \\ \text{all possible } \{A_{\alpha}\} \\ \sum_{\alpha=1}^P Q_{\alpha} = K + 1 ; (P \leq K + 1)}} \frac{1}{Q_1! \cdots Q_P!} \frac{1}{(A_1 + 3)^{Q_1} \cdots (A_P + 3)^{Q_P}} , \end{aligned} \quad (162)$$

where use has been made of

$$\sum_{\beta=1}^P Q_{\beta} (A_{\beta} + 3) = 2(L + K) + 1 , \quad (163)$$

obtained from $\sum_{\beta=1}^P Q_{\beta} A_{\beta} = 2L - K - 2$ and $\sum_{\alpha=1}^P Q_{\alpha} = K + 1$. The relation (159) has been proved. $\square$

Next we show the relation (54) directly: to this end, prove a general formula

$$\int_B^A dt g(t) e^{-Nf(t)} \stackrel{1/N}{=} \int_B^A dt e^{-Nf(t)} . \quad g(t) = -f'(t) + 1 , \quad (164)$$

(In our case, $f(t) \equiv t - \ln t$, $g(t) \equiv 1/t$: $B \equiv 0$, $A \equiv \infty$.) LHS of (164) becomes

$$\text{LHS of (164)} = \int_B^A dt e^{-Nf(t)} - \int_B^A dt f'(t) e^{-Nf(t)} , \quad (165)$$

whose second term is integrable to yield

$$\int_B^A dt f'(t) e^{-Nf(t)} = \int_{f(B)}^{f(A)} df e^{-Nf} = -\frac{1}{N} \left[e^{-Nf(A)} - e^{-Nf(B)} \right] , \quad (166)$$

implying no $1/N$ terms for any value of $Nf(A)$, $Nf(B)$. We have proven (164). $\square$

In our case, from $A = \infty$, $B = 0$, that is, from $f(\infty) = f(0) = \infty$,

$$\int_0^{\infty} dt e^{-N(t - \ln t)} = \int_0^{\infty} dt \frac{1}{t} e^{-N(t - \ln t)} , \quad (167)$$

which designates $\Gamma(N + 1) = N\Gamma(N)$ by multiplying both sides by N^N .

C Calculation of $\mathcal{P}_0^{(\hat{i})} \sim \mathcal{P}_1^{(3)}$ and $\mathcal{L}_0^{(\hat{i})} \sim \mathcal{L}_1^{(3)}$ in (124) and (125)

First expand $F^{(i)}(r); (r = 1, 2, 3)$, (118) (119), such that

$$F^{(i)}(1) \equiv \exp \left[-\frac{N \left(\Omega_c^{(i)} - \omega \right)^2}{2\lambda^2} \right] = F_0^{(i)}(1) \left[1 + \frac{F_1^{(i)}(1)}{N} + \frac{F_2^{(i)}(1)}{N^2} \right], \quad (168)$$

with

$$F_0^{(i)}(1) \equiv \exp \left[-\frac{N \left(\Omega_0^{(i)} - \omega \right)^2}{2\lambda^2} - \frac{\left(\Omega_0^{(i)} - \omega \right) \Omega_1^{(i)}}{\lambda^2} \right], \quad (169)$$

$$F_1^{(i)}(1) \equiv -\frac{\left[\left(\Omega_1^{(i)} \right)^2 + 2 \left(\Omega_0^{(i)} - \omega \right) \Omega_2^{(i)} \right]}{2\lambda^2}, \quad F_2^{(i)}(1) \equiv \frac{\left(F_1^{(i)}(1) \right)^2}{2} - \frac{\Omega_1^{(i)} \Omega_2^{(i)}}{\lambda^2}. \quad (170)$$

$$F^{(i)}(2) \equiv \left(\Omega_c^{(i)} \right)^N = F_0^{(i)}(2) \left[1 + \frac{F_1^{(i)}(2)}{N} + \frac{F_2^{(i)}(2)}{N^2} \right], \quad (171)$$

with

$$F_0^{(i)}(2) \equiv \left(\Omega_0^{(i)} \right)^N \exp \left[\frac{\Omega_1^{(i)}}{\Omega_0^{(i)}} \right], \quad F_1^{(i)}(2) \equiv \frac{\Omega_2^{(i)}}{\Omega_0^{(i)}} - \frac{1}{2} \left(\frac{\Omega_1^{(i)}}{\Omega_0^{(i)}} \right)^2, \quad (172)$$

$$F_2^{(i)}(2) \equiv \frac{\left(F_1^{(i)}(2) \right)^2}{2} - \frac{\Omega_1^{(i)} \Omega_2^{(i)}}{\left(\Omega_0^{(i)} \right)^2} + \frac{1}{3} \left(\frac{\Omega_1^{(i)}}{\Omega_0^{(i)}} \right)^3. \quad (173)$$

(Note that there is no need for $F_2^{(3)}(r); (r = 1, 2)$.)

$$F^{(i)}(3) \equiv \left[1 + \left(\frac{\lambda}{\Omega_c^{(i)}} \right)^2 + \frac{1}{N} \left(\frac{\lambda}{\Omega_c^{(i)} + \delta\omega} \right)^2 \right]^{-1/2} = F_0^{(i)}(3) \left[1 + \frac{F_1^{(i)}(3)}{N} + \frac{F_2^{(i)}(3)}{N^2} \right], \quad (174)$$

with

$$F_0^{(i)}(3) \equiv \begin{cases} \frac{\epsilon \left(\Omega_0^{(\hat{i})} \right) \Omega_0^{(\hat{i})}}{\sqrt{\left(\Omega_0^{(\hat{i})} \right)^2 + \lambda^2}}, \\ \frac{\epsilon \left(\Omega_1^{(3)} \right) \Omega_1^{(3)}}{\sqrt{N} \lambda} \end{cases}, \quad (175)$$

$$F_1^{(i)}(3) \equiv \begin{cases} \frac{\lambda^2}{\left(\Omega_0^{(\hat{i})} \right)^2 + \lambda^2} \left[\frac{\Omega_1^{(\hat{i})}}{\Omega_0^{(\hat{i})}} - \frac{1}{2} \left(\frac{\Omega_0^{(\hat{i})}}{\Omega_0^{(\hat{i})} + \delta\omega} \right)^2 \right] \\ \frac{\Omega_2^{(3)}}{\Omega_1^{(3)}} - \frac{\left(\Omega_1^{(3)} \right)^2}{2\lambda^2} - \frac{1}{2} \left(\frac{\Omega_1^{(3)}}{\Omega_0^{(3)}} \right)^2 \end{cases}. \quad (176)$$

$$F_2^{(\hat{i})}(3) \equiv \frac{3 \left(F_1^{(\hat{i})}(3) \right)^2}{2} + \frac{\lambda^2}{\left(\Omega_0^{(\hat{i})} \right)^2 + \lambda^2} \left[\frac{\Omega_2^{(\hat{i})}}{\Omega_0^{(\hat{i})}} + \frac{\left(\Omega_0^{(\hat{i})} \right)^2 \Omega_1^{(\hat{i})}}{\left(\Omega_0^{(\hat{i})} + \delta\omega \right)^3} - \frac{3}{2} \left(\frac{\Omega_1^{(\hat{i})}}{\Omega_0^{(\hat{i})}} \right)^2 \right]. \quad (177)$$

(Again no need for $F_2^{(3)}(3)$.) Further $F^{(i)}(4)$, (121), for $i = 1, 2$ reads

$$F^{(\hat{i})}(4) = F_0^{(\hat{i})}(4) \left[1 + \frac{F_1^{(\hat{i})}(4)}{N} \right], \quad (178)$$

with

$$F_0^{(\hat{i})}(4) \equiv \left(\Omega_0^{(\hat{i})} + \delta\omega \right)^2 \left(\left(\Omega_0^{(\hat{i})} \right)^2 + \lambda^2 \right); \quad (179)$$

$$F_1^{(\hat{i})}(4) \equiv \frac{2\Omega_1^{(\hat{i})}}{\Omega_0^{(\hat{i})} + \delta\omega} + \frac{2\Omega_0^{(\hat{i})}\Omega_1^{(\hat{i})}}{\left(\Omega_0^{(\hat{i})} \right)^2 + \lambda^2} + \frac{\lambda^2 \left(\Omega_0^{(\hat{i})} \right)^2}{\left(\Omega_0^{(\hat{i})} + \delta\omega \right)^2 \left[\left(\Omega_0^{(\hat{i})} \right)^2 + \lambda^2 \right]}. \quad (180)$$

Meanwhile for $i = 3$

$$NF^{(3)}(4) = F_0^{(3)}(4) \left[1 + \frac{F_1^{(3)}(4)}{N} \right], \quad (181)$$

with

$$F_0^{(3)}(4) \equiv \lambda^2 \left(\Omega_0^{(3)} \right)^2, \quad (182)$$

$$F_1^{(3)}(4) \equiv 2 \frac{\Omega_1^{(3)}}{\Omega_0^{(3)}} + \left(\frac{\Omega_1^{(3)}}{\Omega_0^{(3)}} \right)^2 + \frac{\left(\Omega_1^{(3)} \right)^2}{\lambda^2}. \quad (183)$$

$F^{(M;\hat{i})}(5)$; ($M = 3, \dots, 6$), (122), for $i = 1, 2$ reads

$$F^{(M;\hat{i})}(5) = F_0^{(M;\hat{i})}(5) \left[1 + \frac{F_1^{(M;\hat{i})}(5)}{N} \right], \quad (184)$$

with

$$F_0^{(M;\hat{i})}(5) \equiv \left(\Omega_0^{(\hat{i})} + \delta\omega \right)^M, \quad F_1^{(M;\hat{i})}(5) \equiv \left(\frac{\Omega_0^{(\hat{i})}}{\Omega_0^{(\hat{i})} + \delta\omega} \right)^M + \frac{M\Omega_1^{(\hat{i})}}{\Omega_0^{(\hat{i})} + \delta\omega}. \quad (185)$$

Then for $i = 3$

$$NF^{(M;3)}(5) = F_0^{(M;3)}(5) \left[1 + \frac{F_1^{(M;3)}(5)}{N} \right], \quad M = 3, \dots, 6, \quad (186)$$

with

$$F_0^{(M;3)}(5) \equiv \left(\Omega_0^{(3)} \right)^M; \quad F_1^{(M;3)}(5) \equiv M \frac{\Omega_1^{(3)}}{\Omega_0^{(3)}}. \quad (187)$$

Therefore the prefactor reads

$$\begin{aligned}\mathcal{P}_0^{(i)} &\equiv F_0^{(i)}(1)F_0^{(i)}(2)F_0^{(i)}(3) \left(\Omega_0^{(i)} + \delta\omega \right) \\ &= \exp \left[-\frac{N \left(\Omega_0^{(i)} - \omega \right)^2}{2\lambda^2} \right] \left(\Omega_0^{(i)} \right)^{N+1} \frac{\epsilon \left(\Omega_0^{(i)} \right)}{\sqrt{\left(\Omega_0^{(i)} \right)^2 + \lambda^2}} \left(\Omega_0^{(i)} + \delta\omega \right),\end{aligned}\quad (188)$$

for $i = 1, 2$ and the leading term of $i = 3$ is

$$\mathcal{P}_1^{(3)} = \exp \left[-\frac{N \left(\Omega_0^{(3)} - \omega \right)^2}{2\lambda^2} \right] \left(\Omega_0^{(3)} \right)^N \frac{\epsilon \left(\Omega_1^{(3)} \right) \left(\Omega_1^{(3)} \right)^2}{e\sqrt{N}\lambda}, \quad (189)$$

where use has been made of the relation

$$-\frac{\left(\Omega_0^{(i)} - \omega \right) \Omega_1^{(i)}}{\lambda^2} + \frac{\Omega_1^{(i)}}{\Omega_0^{(i)}} = \begin{cases} 0 & i = 1, 2 \\ -1 & i = 3 \end{cases}, \quad (190)$$

in view of (113) and the gap equation (83) for $i = 1, 2$ or (114) with (115) for $i = 3$. Then

$$\mathcal{P}_1^{(i)} = \sum_{r=1}^3 F_1^{(i)}(r) + \frac{\Omega_1^{(i)}}{\Omega_0^{(i)} + \delta\omega}, \quad (191)$$

$$\mathcal{P}_2^{(3)} = \sum_{r=1}^3 F_1^{(3)}(r) + \frac{\Omega_2^{(3)}}{\Omega_1^{(3)}}, \quad (192)$$

and

$$\mathcal{P}_2^{(i)} = \sum_{r=1}^3 F_2^{(i)}(r) + \frac{\Omega_2^{(i)}}{\Omega_0^{(i)} + \delta\omega} + \sum_{r' > r=1}^3 F_1^{(i)}(r') F_1^{(i)}(r) + \frac{\Omega_1^{(i)}}{\Omega_0^{(i)} + \delta\omega} \sum_{r=1}^3 F_1^{(i)}(r). \quad (193)$$

Now calculate terms in the loop factor (120) up to the 3-loop approximation:

$$\begin{aligned}\mathcal{L}^{(i)} &= 1 + \frac{\lambda^2}{NF^{(i)}(4)} \left[-\frac{3\lambda^2}{4} R^{(4;i)} + \frac{5\lambda^4}{6} \left(R^{(3;i)} \right)^2 \right] \\ &\quad + \frac{\lambda^4}{\left(NF^{(i)}(4) \right)^2} \left[-\frac{5\lambda^2}{2} R^{(6;i)} + 7\lambda^4 R^{(3;i)} R^{(5;i)} + \frac{105\lambda^4}{32} \left(R^{(4;i)} \right)^2 \right. \\ &\quad \left. - \frac{105\lambda^6}{8} \left(R^{(3;i)} \right)^2 R^{(4;i)} + \frac{385\lambda^8}{72} \left(R^{(3;i)} \right)^4 \right],\end{aligned}\quad (194)$$

where we have introduced

$$R^{(M;i)} \equiv \frac{F^{(M;i)}(5)}{F^{(i)}(4)} = R_0^{(M;i)} \left[1 + \frac{R_1^{(M;i)}}{N} \right] + O\left(\frac{1}{N^2} \right), \quad (195)$$

whose expansion coefficients are

$$R_0^{(M;i)} \equiv \frac{F_0^{(M;i)}(5)}{F_0^{(i)}(4)} = \begin{cases} \frac{\left(\Omega_0^{(i)} + \delta\omega \right)^{M-2}}{\left(\Omega_0^{(i)} \right)^2 + \lambda^2} \\ \frac{\left(\Omega_0^{(3)} \right)^{M-2}}{\lambda^2} \end{cases}, \quad (196)$$

$$\begin{aligned}
R_1^{(M;i)} &\equiv F_1^{(M;i)}(5) - F_1^{(i)}(4) \\
&= \begin{cases} \left(\frac{\Omega_0^{(i)}}{\Omega_0^{(i)} + \delta\omega} \right)^M + \frac{(M-2)\Omega_1^{(i)}}{\Omega_0^{(i)} + \delta\omega} - \frac{2\Omega_0^{(i)}\Omega_1^{(i)}}{\left(\Omega_0^{(i)}\right)^2 + \lambda^2} - \frac{\lambda^2 \left(\Omega_0^{(i)}\right)^2}{\left(\Omega_0^{(i)} + \delta\omega\right)^2 \left[\left(\Omega_0^{(i)}\right)^2 + \lambda^2\right]} \\ (M-2)\frac{\Omega_1^{(3)}}{\Omega_0^{(3)}} - \left(\frac{\Omega_1^{(3)}}{\Omega_0^{(3)}}\right)^2 - \frac{\left(\Omega_1^{(3)}\right)^2}{\lambda^2} \end{cases} .(197)
\end{aligned}$$

With the aid of these, $i = 1, 2$ part is given by

$$\mathcal{L}_0^{(i)} = 1 ; \quad (198)$$

$$\begin{aligned}
\mathcal{L}_1^{(i)} &= \frac{\lambda^2}{\left(\Omega_0^{(i)} + \delta\omega\right)^2 \left[\left(\Omega_0^{(i)}\right)^2 + \lambda^2\right]} \left[-\frac{3\lambda^2}{4} R_0^{(4;i)} + \frac{5\lambda^4}{6} \left(R_0^{(3;i)}\right)^2 \right] \\
&= -\frac{3\lambda^4}{4 \left[\left(\Omega_0^{(i)}\right)^2 + \lambda^2\right]^2} + \frac{5\lambda^6}{6 \left[\left(\Omega_0^{(i)}\right)^2 + \lambda^2\right]^3} , \quad (199)
\end{aligned}$$

$$\begin{aligned}
\mathcal{L}_2^{(i)} &= \frac{\lambda^4}{\left(\Omega_0^{(i)} + \delta\omega\right)^4 \left[\left(\Omega_0^{(i)}\right)^2 + \lambda^2\right]^2} \left[-\frac{5\lambda^2}{2} R_0^{(6;i)} + 7\lambda^4 R_0^{(3;i)} R_0^{(5;i)} + \frac{105\lambda^4}{32} \left(R_0^{(4;i)}\right)^2 \right. \\
&\quad \left. - \frac{105\lambda^6}{8} \left(R_0^{(3;i)}\right)^2 R_0^{(4;i)} + \frac{385\lambda^8}{72} \left(R_0^{(3;i)}\right)^4 \right] \\
&\quad - \mathcal{L}_1^{(i)} F_1^{(i)}(4) - \frac{3\lambda^4}{4} \frac{R_0^{(4;i)} R_1^{(4;i)}}{F_0^{(i)}(4)} + \frac{5\lambda^6}{3} \frac{\left(R_0^{(3;i)}\right)^2 R_1^{(3;i)}}{F_0^{(i)}(4)} \\
&= -\frac{5}{2} \frac{\lambda^6}{\left[\left(\Omega_0^{(i)}\right)^2 + \lambda^2\right]^3} + \frac{329}{32} \frac{\lambda^8}{\left[\left(\Omega_0^{(i)}\right)^2 + \lambda^2\right]^4} - \frac{105}{8} \frac{\lambda^{10}}{\left[\left(\Omega_0^{(i)}\right)^2 + \lambda^2\right]^5} \\
&\quad + \frac{385}{72} \frac{\lambda^{12}}{\left[\left(\Omega_0^{(i)}\right)^2 + \lambda^2\right]^6} - \mathcal{L}_1^{(i)} F_1^{(i)}(4) - \frac{3\lambda^4}{4} \frac{R_0^{(4;i)} R_1^{(4;i)}}{F_0^{(i)}(4)} + \frac{5\lambda^6}{3} \frac{\left(R_0^{(3;i)}\right)^2 R_1^{(3;i)}}{F_0^{(i)}(4)} . \quad (200)
\end{aligned}$$

For $i = 3$, in view of (181), a naive $1/N$ expansion is broken down; all factors in front of the loop expansion, (120), behave $O(1)$. Nevertheless we take up to the terms of $O(1/\{NF^{(3)}(4)\})$ under the 2-loop approximation, and of $O(1/\{NF^{(3)}(4)\}^2)$ under 3-loop so that

$$\mathcal{L}_0^{(3)} \Big|_{2\text{-loop}} = 1 + \frac{\lambda^2}{F_0^{(3)}(4)} \left[-\frac{3\lambda^2}{4} R_0^{(4;3)} + \frac{5\lambda^4}{6} \left(R_0^{(3;3)}\right)^2 \right] = \frac{13}{12} , \quad (201)$$

with the use of (182) and (196), and for $O(1/N)$ terms we obtain

$$\begin{aligned} \mathcal{L}_1^{(3)} \Big|_{2\text{-loop}} &= \frac{\lambda^2}{F_0^{(3)}(4)} \left[\left\{ -\frac{3\lambda^2}{4} R_0^{(4;3)} + \frac{5\lambda^4}{6} \left(R_0^{(3;3)} \right)^2 \right\} \left(-F_1^{(3)}(4) \right) + \left\{ -\frac{3\lambda^2}{4} R_0^{(4;3)} R_1^{(4;3)} \right. \right. \\ &\quad \left. \left. + \frac{5\lambda^4}{3} \left(R_0^{(3;3)} \right)^2 R_1^{(3;3)} \right\} \right] = -\frac{1}{12} F_1^{(3)}(4) - \frac{3}{4} R_1^{(4;3)} + \frac{5}{3} R_1^{(3;3)} \\ &= - \left[\left(\frac{\Omega_1^{(3)}}{\Omega_0^{(3)}} \right)^2 + \frac{\left(\Omega_1^{(3)} \right)^2}{\lambda^2} \right], \end{aligned} \quad (202)$$

where use has been made of (182), (183), (196), and (197).

In a similar manner, we obtain the $O(1)$ term in 3-loop, such that

$$\begin{aligned} \mathcal{L}_0^{(3)} \Big|_{3\text{-loop}} &= \mathcal{L}_0^{(3)} \Big|_{2\text{-loop}} + \frac{\lambda^4}{\left(F_0^{(3)}(4) \right)^2} \left[-\frac{5\lambda^2}{2} R_0^{(6;3)} + 7\lambda^4 R_0^{(3;3)} R_0^{(5;3)} + \frac{105\lambda^4}{32} \left(R_0^{(4;3)} \right)^2 \right. \\ &\quad \left. - \frac{105\lambda^6}{8} \left(R_0^{(3;3)} \right)^2 R_0^{(4;3)} + \frac{385\lambda^8}{72} \left(R_0^{(3;3)} \right)^4 \right] = \frac{313}{288}, \end{aligned} \quad (203)$$

and for $O(1/N)$ terms

$$\begin{aligned} \mathcal{L}_1^{(3)} \Big|_{3\text{-loop}} &= \mathcal{L}_1^{(3)} \Big|_{2\text{-loop}} + \frac{\lambda^4}{\left(F_0^{(3)}(4) \right)^2} \left[\left\{ -\frac{5\lambda^2}{2} R_0^{(6;3)} + 7\lambda^4 R_0^{(3;3)} R_0^{(5;3)} + \frac{105\lambda^4}{32} \left(R_0^{(4;3)} \right)^2 \right. \right. \\ &\quad \left. \left. - \frac{105\lambda^6}{8} \left(R_0^{(3;3)} \right)^2 R_0^{(4;3)} + \frac{385\lambda^8}{72} \left(R_0^{(3;3)} \right)^4 \right\} \left(-2F_1^{(3)}(4) \right) \right. \\ &\quad \left. + \left\{ -\frac{5\lambda^2}{2} R_0^{(6;3)} R_1^{(6;3)} + 7\lambda^4 R_0^{(3;3)} R_0^{(5;3)} \left(R_1^{(3;3)} + R_1^{(5;3)} \right) + \frac{105\lambda^4}{16} \left(R_0^{(4;3)} \right)^2 R_1^{(4;3)} \right. \right. \\ &\quad \left. \left. - \frac{105\lambda^6}{8} \left(R_0^{(3;3)} \right)^2 R_0^{(4;3)} \left(2R_1^{(3;3)} + R_1^{(4;3)} \right) + \frac{385\lambda^8}{18} \left(R_0^{(3;3)} \right)^4 R_1^{(3;3)} \right\} \right] \\ &= -\frac{13}{144} F_1^{(3)}(4) + \frac{137}{36} R_1^{(3;3)} - \frac{117}{16} R_1^{(4;3)} + 7R_1^{(5;3)} - \frac{5}{2} R_1^{(6;3)} \\ &= -\frac{13}{12} \left[\left(\frac{\Omega_1^{(3)}}{\Omega_0^{(3)}} \right)^2 + \frac{\left(\Omega_1^{(3)} \right)^2}{\lambda^2} \right]. \end{aligned} \quad (204)$$

References

- [1] T. Kashiwa, Y. Ohnuki and M. Suzuki, *Path Integral Methods* (Clarendon Press · Oxford, 1997) p.162.
- [2] D.Gross and A. Neveu, Phys. Rev. **D10**, 3235 (1974).
T. Kugo, Prog. Theor. Phys. **55**, 2032 (1976).
K. Kikkawa, Prog. Theor. Phys. **56**, 947 (1976).
- [3] E. Fradkin, *Field Theories of Condensed Matter System* (Addison-Wesley, New York, 1991) p.327.

- [4] J. Zinn-Justin, *Quantum Field Theory and Critical Phenomena* (Clarendon Press · Oxford, 1989) Chap.5.
Also sec.2.3 in ref.[1].
- [5] R. P. Feynman and A. R. Hibbs, *Quantum Mechanics and Path Integrals* (McGraw-Hill, New York,1965) Chap.11.
H. Kleinert, *Path Integrals* (World Scientific, Singapore, 1995) Chap. 5.
- [6] A. Okopińska, Phys. Rev. **D 35** (1987) 1835; **D 36** (1987) 2415.
I. R. C. Buckley, A. Duncan, and H. F. Jones, Phys. Rev. **D 47** (1993) 2554.
A. Duncan and H. F. Jones, Phys. Rev. **D 47** (1993) 2560.
- [7] T. Kashiwa, Phys. Rev. **D59**, 085002 (1999).
- [8] T. Kashiwa and T.Sakaguchi, Prog. Theor. Phys. **110** , 589 (2003)
- [9] T. Kashiwa and T.Sakaguchi, J.Phys.B At.Mol.Opt.Phys. **39**, S787 (2006).
- [10] I.S. Gradshteyn and I.M. Ryzhik, *Tables of Integrals Series, and Product: 6th Ed.*(Academic Press, 2000) p.886, 8.327
- [11] S. Moriguchi, KDUdagawa and S.Hitotsumatsu, *Tables of Mathematical Formula III* (In Japanese) p. 5 (Iwanami.1987)
- [12] function.wolfram.com, *Gamma function:Series representations*,
<http://functions.wolfram.com/GammaBetaErf/Gamma/06/02/0002/>.
- [13] E.T. Copson, *Asymptotic Expansions* (Cambridge 1965) Chap.6
- [14] Y. Nambu and G. Jona-Lasinio, Phys. Rev. **122**, 345 (1961).
M. Ishi-i, T. Kashiwa, and N. Tanimura, Prog. Theor. Phys. **100**, 353 (1998).
M. Ishi-i, T. Kashiwa, and N. Tanimura, Phys. Rev. **D65**, 065025 (2002).
T. Kashiwa and T. Sakaguchi, Phys. Rev. **D 68** 065002 (2003).